arXiv:math/0702451v4 [math.AC] 16 Jul 2008

Commutative rings with toroidal zero-divisor graphs

Hung-Jen Chiang-Hsieh* and Hsin-Ju Wang†
Department of Mathematics,
National Chung Cheng University,
Chiayi 621, Taiwan

Neal O. Smith‡
Department of Mathematics,
Augusta State University,
Augusta GA 30904

Abstract

Let R be a commutative ring and let $\Gamma(R)$ denote its zero-divisor graph. We investigate the genus number of the compact Riemann surface in which $\Gamma(R)$ can be embedded and explicitly determine all finite commutative rings R (up to isomorphism) such that $\Gamma(R)$ is either toroidal or planar.

Introduction

We assume that all rings are commutative with identity. For a ring R, the zero-divisor graph of R, denoted by $\Gamma(R)$, is the simple graph whose vertex set consists of all nonzero zero-divisors of R. Two distinct vertices are joined by an edge if and only if the product of the vertices is 0. Therefore $\Gamma(R) = \emptyset$ if and only if R is an integral domain. This definition was introduced by Anderson and Livingston in [6]. Recently, this subject has been extensively studied in [2], [3], [4], [5], [7], [8], [12], [13], [15], [18], [19], and [20].

There are many known results concerning zero-divisor graphs. Anderson and Livingston showed in [6] that $\Gamma(R)$ is always connected and R is a finite ring or an integral domain if and only if $\Gamma(R)$ is finite. Mulay [15] showed that if $\Gamma(R)$ contains a cycle, then $\Gamma(R)$ contains a 3-cycle or a 4-cycle. Anderson, Frazier, Lauve and Livingston showed in [4] that if R and S are finite reduced rings which are not fields, then $R \simeq S$ if and only if $\Gamma(R) \simeq \Gamma(S)$.

The main objective of topological graph theory is to embed a graph into a surface. Simply stated, that is to draw a graph on a surface so that no two edges cross one another. The simplest case of this problem is when the surface in question is the plane; if a graph can be embedded in the plane, we say the graph is *planar*. There are many papers where planarity of zero-divisor graphs has been discussed. In [4], Anderson et al. the authors determined when $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$ and $\mathbb{Z}_n[X]/(X^m)$ have planar zero

*e-mail: hchiang@math.ccu.edu.tw (corresponding author)
†e-mail: hjwang@math.ccu.edu.tw
‡e-mail: nsmith12@aug.edu
2000 Mathematics Subject Classification. Primary 13A99, 05C10, 13M99.

divisor graphs and posed the general question as to which finite rings R have $\Gamma(R)$ planar. It was shown in [2] that if R is a finite local ring such that $\Gamma(R)$ has at least 33 vertices, then $\Gamma(R)$ is not planar. In that paper, Akbari et al. conjectured that for any local ring of cardinality 32 which is not a field, $\Gamma(R)$ is not planar.

This conjecture was proved independently in two papers. In [18], Smith proved that if R is a finite, commutative local ring (not a field) with cardinality 28 or greater, then $\Gamma(R)$ is not planar. Also in that paper the author classifies precisely those finite commutative rings R for which $\Gamma(R)$ is planar. Some of the methods used in that paper are similar in spirit to some of the arguments in section three of this paper; thus, we recover this listing of planar graphs as we work toward our main theorem. We also refer the readers to [7], in which Belshoff and Chapman independently obtain some of the same results from [18], and to [16], in which Redmond lists all zero-divisor graphs up to 14 vertices where some of them are among the same results as above.

Akbari's conjecture was also verified independently in [20]. Moreover, in the same paper the author also found all finite rings of the form $\mathbb{Z}_{p_1^{\alpha_1}} \times \cdots \times \mathbb{Z}_{p_n^{\alpha_n}}$ or of the form $\mathbb{Z}_n[x]/(x^m)$ whose zero-divisor graphs are planar or can be embedded into a torus. This motivates the work in this paper.

To find all finite rings R such that $\Gamma(R)$ has genus at most one is the goal of this paper. Since a finite ring is Artinian, it is a direct product of local Artinian rings. Thus, we first consider the case of finite local rings. To motivate the main theorems in section 3, we first discuss the genera of local rings under some specific assumptions in section 2. Using the Euler characteristic formula and a technique of deletion and insertion, we are able to successfully exclude some cases of higher genus.

In section 3, we consider case by case those local rings $(R, \mathfrak{m})$ with $|R/\mathfrak{m}| \leq 8$. From [20, Theorem 3.6], we know that these are all the cases of interest. By [20, Lemma 3.1], if a finite ring R has $|Spec(R)| \geq 5$ then $\gamma(\Gamma(R)) \geq 2$. According to this, it suffices to look for the finite rings with at most 4 maximal ideals. We obtain a complete list of all finite rings whose zero-divisor graphs have genera at most one in this section and summarize them in four tables at the very end of this paper. Finally, we do a similar analysis in the case where R decomposes as a product of local rings.

We should mention that as this paper was being submitted, the authors became aware of a similar work. In [21], Wickham independently proved some of the same results that appear in section three of this paper.

1 Preliminaries

In this section we briefly recap some notation, terminology, and basic results from [11], [17] and [20].

A simple graph G is an ordered pair of disjoint sets (V, E), such that $V = V(G)$ is the set of vertices of G and $E = E(G)$ is the set of edges of G. For $v \in V$, the *degree* of v, denoted by $deg(v)$, is the number of edges of G incident to v. If $V' \subseteq V(G)$, we define $G - V'$ to be the subgraph of G obtained by deleting the vertices in V' and all incident edges. Similarly, if $E' \subseteq E(G)$, then $G - E'$ is the subgraph of G obtained by deleting the edges in E'. For a graph G, let $\widetilde{G}$ denote the subgraph $G - V'$ where $V' = \{ v \in V \mid deg(v) = 1\}$. We call this graph the *reduction* of G.

A graph in which each pair of distinct vertices is joined by an edge is called a *complete graph*. We use K_n to denote a complete graph with n vertices. A *bipartite graph* G is a graph whose vertex set $V(G)$ can be partitioned into two subsets V_1 and V_2. The edge set of such a graph consists of precisely those edges which join vertices in V_1 to vertices of V_2. In particular, if $E(G)$ consists of all possible such edges, then G called a *complete bipartite graph* and denoted by the symbol $K_{m,n}$ where $|V_1| = m$ and $|V_2| = n$.

By a *surface*, we mean a two dimensional real manifold, that is a topological space such that each

point has a neighborhood homeomorphic to the open disc. It is well-known that every orientable compact surface is homeomorphic to a sphere with g handles. This number g is called the genus of the surface. For example, the genus of a sphere is 0 and the genus of a torus is 1. A simple graph which can be drawn without crossings on a surface of genus g but not on one of genus $g-1$, is called a graph of genus g. We say a *planar* graph is a graph of genus 0 and a *toroidal* graph is a graph of genus 1. We use $\gamma(G)$ to denote the genus of a graph G. The following two results (see [11, p.118]) about the genus of a complete graph and a complete bipartite graph will be very useful in the subsequent sections.

Lemma 1.1. *Let $n \geq 3$. Then $\gamma(K_n) = \{\frac{1}{12}(n-3)(n-4)\}$, where $\{x\}$ is the least integer that is greater than or equal to x. In particular, $\gamma(K_n) = 1$ if $n = 5, 6,$ or 7.*

Lemma 1.2. *$\gamma(K_{m,n}) = \{\frac{1}{4}(m-2)(n-2)\}$, where $\{x\}$ is the least integer that is greater than or equal to x. In particular, $\gamma(K_{4,4}) = \gamma(K_{3,n}) = 1$ if $n = 3, 4, 5,$ or 6.*

Suppose a connected graph G is drawn on an orientable compact surface S_g and let $\#V_G, \#E_G$, and $\#F_G$ denote the number of vertices, edges, and faces of G respectively. The well known Euler characteristic formula states that $\#V_G - \#E_G + \#F_G = 2 - 2g$, where g is the genus of S_g.

We end this section with two remarks.

Remark 1.3. $\gamma(H) \leq \gamma(G)$ for all subgraphs H of G; and $\gamma(\widetilde{G}) = \gamma(G)$, where $\widetilde{G}$ is the reduction of G.

Remark 1.4. The bipartite graph $K_{3,6}$ has $v = 9$ vertices and $e = 18$ edges. By Lemma 1.2, we see that $\gamma(K_{3,6}) = 1$. Therefore, from the Euler characteristic formula there are $f = 9$ faces when drawing $K_{3,6}$ without crossings on a torus. We note that the boundary of each face F_i of $K_{3,6}$ is an even cycle with length $e_i \geq 4$. It follows from the inequality $2e = \sum_{i=1}^{f} e_i \geq 4f$ that $e_i = 4$ for each i. That is, all face boundaries are 4-cycles. Moreover, any two faces in $K_{3,6}$ have at most one boundary edge in common.

2 Genera of some special rings

In order to simplify the proof of our main result in the next section, we now discuss several special rings with $\gamma(\Gamma(R)) \geq 2$ or $\gamma(\Gamma(R)) = 1$. In the sequel, if R is a ring then $Z(R)$ will denote the set of its zero-divisors and we define $Z(R)^* = Z(R) - \{0\}$.

Proposition 2.1. *Let $(R, \mathfrak{m})$ be a local ring with $|R| = 32$. If $|R/\mathfrak{m}| = 2$ and $|\mathfrak{m}^2| = 4$, then $\gamma(\Gamma(R)) \geq 2$.*

Proof. We have that $|\mathfrak{m}| = 16$ and $dim_{R/\mathfrak{m}}\mathfrak{m}/\mathfrak{m}^2 = 2$. Since $\mathfrak{m}^2 \neq \{0\}$, by Nakayama's lemma [1, Prop. 2.6] we have $\mathfrak{m}^3 \subsetneq \mathfrak{m}^2$. Thus $\mathfrak{m}^3 = \{0\}$ or $|\mathfrak{m}^3| = 2$. If $\mathfrak{m}^3 = \{0\}$, then $|\mathfrak{m} - \mathfrak{m}^2| = 12$ and $|\mathfrak{m}^2 - \mathfrak{m}^3| = 3$. This implies that $K_{3,12} \subseteq \Gamma(R)$ and therefore $3 = \gamma(K_{3,12}) \leq \gamma(\Gamma(R))$ by Lemma 1.2 and Remark 1.3. Hence, we may assume $|\mathfrak{m}^3| = 2$, so that $\mathfrak{m}^4 = \{0\}$ and $dim_{R/\mathfrak{m}}\mathfrak{m}^2/\mathfrak{m}^3 = dim_{R/\mathfrak{m}}\mathfrak{m}^3/\mathfrak{m}^4 = 1$.

Since $dim_{R/\mathfrak{m}}\mathfrak{m}/\mathfrak{m}^2 = 2$, it follows from [1, Prop.2.8] that $\mathfrak{m}$ can be generated by two elements. Write $\mathfrak{m} = (x, y)$. Assume first that $x^3 = y^3 = 0$. If $x^2y \neq 0$, then $x^2, xy \notin \mathfrak{m}^3$, so that $\{x^2\}$ and $\{xy\}$ are both bases for $\mathfrak{m}^2/\mathfrak{m}^3$. It then follows that $xy - x^2 \in \mathfrak{m}^3$ as $|R/\mathfrak{m}| = 2$. Therefore, we have $x^2y - x^3 = 0$ and $x^2y = 0$, a contradiction. Thus, $x^2y = 0$ and $xy^2 = 0$. If $x^3 = y^3 = 0$ we have $\mathfrak{m}^3 = \{0\}$, a contradiction. Consequently, we conclude that either $x^3 \neq 0$ or $y^3 \neq 0$. We may assume without loss that $x^3 \neq 0$. Therefore, $\{x^3\}$ is a basis for $\mathfrak{m}^3/\mathfrak{m}^4$ and $\{x^2\}$ is a basis for $\mathfrak{m}^2/\mathfrak{m}^3$. If $xy \notin \mathfrak{m}^3$, then $xy - x^2 \in \mathfrak{m}^3$ as

$dim_{R/\mathfrak{m}}\mathfrak{m}^2/\mathfrak{m}^3 = 1$ and $|R/\mathfrak{m}| = 2$, so that $x(y-x) \in \mathfrak{m}^3$. Therefore, we may replace y by $y-x$ and assume that $xy \in \mathfrak{m}^3$. Moreover, if $xy \neq 0$, then $xy - x^3 = 0$ as $dim_{R/\mathfrak{m}}\mathfrak{m}^3/\mathfrak{m}^4 = 1$ and $|R/\mathfrak{m}| = 2$, so that $x(y-x^2) = 0$. Hence we may replace y by $y-x^2$ and assume that $xy = 0$. We observe that $y^2 \in \mathfrak{m}^3$. If not, then $y^2 - x^2 \in \mathfrak{m}^3$, so that $xy^2 - x^3 = 0$. It then follows that $x^3 = 0$, a contradiction. Now, we have two cases to discuss, as either $y^2 = 0$ or $y^2 = x^3$.

Case 1 : $y^2 = 0$. In this case, let $u_1 = x^3$, $u_2 = y$, $u_3 = y + x^3$, $v_1 = x$, $v_2 = x + x^2$, $v_3 = x + x^3$, $v_4 = x + x^2 + x^3$, $v_5 = x^2$, $v_6 = x^2 + x^3$ and $v_7 = x + y$. Then $u_i \cdot v_j = 0$ for every i, j. Therefore $K_{3,7} \subseteq \Gamma(R)$ and it follows that $\gamma(\Gamma(R)) \geq \gamma(K_{3,7}) = 2$.

Case 2 : $y^2 = x^3$. In this case, let $u_1 = x^3$, $u_2 = y$, $u_3 = y + x^3$, $v_1 = x$, $v_2 = x + x^2$, $v_3 = x + x^3$, $v_4 = x + x^2 + x^3$, $v_5 = x^2$, $v_6 = x^2 + x^3$, $w_1 = x + y$, $w_2 = x + y + x^2$, $w_3 = x + y + x^3$, $w_4 = x + y + x^2 + x^3$, $w_5 = y + x^2$ and $w_6 = y + x^2 + x^3$. Observe that $u_i \cdot v_j = 0$ for every i, j, so that $K_{3,6} \subseteq \Gamma(R)$. Therefore $\gamma(\Gamma(R)) \geq \gamma(K_{3,6}) = 1$. Write $G = \Gamma(R)$, $G' = G - \{u_1u_2, u_1u_3, u_1w_5, u_1w_6\}^*$, and $G'' = G' - \{w_1, \ldots, w_6\}$. It is then easy to see that $G'' \simeq K_{3,6}$. Next, we proceed to prove $\gamma(G) \geq 2$ by a deletion and insertion argument.

Suppose that $\gamma(G) = 1$. Since $1 = \gamma(G'') \leq \gamma(G') \leq \gamma(G)$, we get $\gamma(G') = 1$. Since

$$V(G') = \{u_1, u_2, u_3, v_1, \ldots, v_6, w_1, \ldots, w_6\}$$

and

$$\begin{aligned} E(G') = \quad & \{\, u_iv_j \,|\, 1 \leq i \leq 3, 1 \leq j \leq 6 \,\} \cup \{\, u_1w_i \,|\, 1 \leq i \leq 4\} \\ & \cup \{\, w_iw_j \,|\, 1 \leq i \leq 4, j = 5, 6\} \cup \{\, v_5w_5, v_5w_6, v_6w_5, v_6w_6 \,\}, \end{aligned}$$

by the Euler characteristic formula there are 19 faces when drawing G' on a torus. Fix a representation of G' and let $\{F'_1, \ldots, F'_{19}\}$ be the set of faces of G' corresponding to this representation. We note that $G'' \simeq K_{3,6}$ and therefore this graph has 9 faces whose boundaries are all 4-cycles (see Remark 1.4). Write $F''_1, \ldots, F''_9$ for the faces of G'' obtained by deleting $w_1, \ldots, w_6$ and all edges incident with $w_1, \ldots, w_6$ from the representation of G'. Then $\{F'_1, \ldots, F'_{19}\}$ can be recovered by inserting $w_1, \ldots, w_6$ and all edges incident with $w_1, \ldots, w_6$ into the representation corresponding to $\{F''_1, \ldots, F''_9\}$. Let F''_{t_i} denote the face of G'' into which w_i is inserted during the recovering process from G'' to G'. We note that $w_iw_j \in E(G')$ for $i = 1, \ldots, 4$ and $j = 5, 6$. Therefore every w_i should be inserted into the same face, say F''_m, of G'' to avoid any crossings, i.e., $t_1 = t_2 = \cdots = t_6 = m$. Moreover, since $u_1w_i \in E(G')$ for $i = 1, \ldots, 4$, u_1 is a vertex of the face F''_m. Write the edges $e_i = u_1w_i$, $e_{i+4} = w_iw_5$ and $e_{i+8} = w_iw_6$ for $i = 1, \ldots, 4$. After inserting $w_1, \ldots, w_5$ and $e_1, \ldots, e_8$ into F''_m we obtain Figure 1 as below. However, it is easy to see from Figure 1 that we can not insert w_6 and $e_9, \ldots, e_{12}$ into F''_m without crossings, a contradiction. Therefore, we may conclude that $\gamma(G) \geq 2$. □



Figure 1

*If $u, v \in R$ such that $uv = 0$ then we write uv as an edge of $\Gamma(R)$.

Proposition 2.2. *Let $(R, \mathfrak{m})$ be a local ring with $|R| = 32$. If $|R/\mathfrak{m}| = 2$ and $|\mathfrak{m}^2| = 2$, then $\gamma(\Gamma(R)) \geq 2$.*

Proof. Note that $|\mathfrak{m}| = 16$, $\mathfrak{m}^3 = \{0\}$, $dim_{R/\mathfrak{m}}\mathfrak{m}/\mathfrak{m}^2 = 3$ and $dim_{R/\mathfrak{m}}\mathfrak{m}^2/\mathfrak{m}^3 = 1$. Therefore, by [1, Prop.2.8], $\mathfrak{m}$ can be generated by three elements, say, $\mathfrak{m} = (x, y, z)$ for some $x, y, z \in \mathfrak{m} - \mathfrak{m}^2$.

First we assume that $x\mathfrak{m} = \{0\}$. Choose $w \in \mathfrak{m}^2 - \{0\}$ and denote $u_1 = x$, $u_2 = w$, $u_3 = x + w$, $v_1 = y$, $v_2 = z$, $v_3 = y + z$, $v_4 = y + w$, $v_5 = z + w$, $v_6 = y + z + w$, and $v_7 = x + y$; then $u_i \cdot v_j = 0$ for every i, j. It follows that $K_{3,7} \subseteq \Gamma(R)$, so that $\gamma(\Gamma(R)) \geq \gamma(K_{3,7}) = 2$. Thus we may assume that $u\mathfrak{m} \neq \{0\}$ for any $u \in \mathfrak{m} - \mathfrak{m}^2$.

Suppose that $x^2 = y^2 = z^2 = 0$. Since $2ab = 0$ for any $a, b \in \mathfrak{m}$ since $\mathfrak{m}^3 = \{0\}$ and $|R/\mathfrak{m}| = 2$, we have that $u^2 = 0$ for all $u \in \mathfrak{m}$. Noting that $|\mathfrak{m}^2| = 2$, we may assume $xy \neq 0$, that is, $\mathfrak{m}^2 = \{xy, 0\}$. Now if $xz \neq 0$, then $xz = xy$, so that $x(z - y) = 0$. Replacing z by $z - y$, we may assume $xz = 0$. Furthermore, if $yz \neq 0$, then $yz = xy$, so that $y(z - x) = 0$. Since $x(z - x) = 0$, we may replace z by $z - x$ and assume that $yz = 0$. However, this implies $z\mathfrak{m} = \{0\}$, which contradicts the assumption that $u\mathfrak{m} \neq \{0\}$ for any $u \in \mathfrak{m} - \mathfrak{m}^2$. Thus, $u^2 \neq 0$ for some $u \in \{x, y, z\}$. After a suitable change of x, y, z, we may assume that $x^2 \neq 0$ and $xy = xz = 0$. There are thus two cases to consider, either $y^2 \neq 0$ or $y^2 = 0$.
Case 1 : $y^2 \neq 0$. In this case, we may further assume that $yz = 0$ as in the previous paragraph. Therefore $z^2 \neq 0$ as $z\mathfrak{m} \neq \{0\}$. Consequently, $x^2 = y^2 = z^2$ and $xy = xz = yz = 0$. Let $u_1 = x^2$, $u_2 = x$, $u_3 = x + x^2$, $v_1 = y$, $v_2 = z$, $v_3 = y + z$, $v_4 = y + x^2$, $v_5 = z + x^2$, $v_6 = y + z + x^2$, $w_1 = x + y$, $w_2 = x + y + x^2$, $w_3 = x + z$, $w_4 = x + z + x^2$, $w_5 = x + y + z$ and $w_6 = x + y + z + x^2$. Then $u_i \cdot v_j = 0$ for every i, j. It follows that $K_{3,6} \subseteq \Gamma(R)$, and therefore $\gamma(\Gamma(R)) \geq \gamma(K_{3,6}) = 1$. Let $G = \Gamma(R)$, $G' = G - \{u_1w_1, u_1w_2, \ldots, u_1w_6, u_1u_2, u_1u_3, v_1v_2, v_1v_5, v_2v_4, v_3v_6\}$, and $G'' = G' - \{w_1, w_2, \ldots, w_6\}$. We then have $G'' \simeq K_{3,6}$.

Suppose that $\gamma(G) = 1$. Since $1 = \gamma(G'') \leq \gamma(G') \leq \gamma(G)$, we have $\gamma(G') = 1$. Since

$$V(G') = \{u_1, u_2, u_3, v_1, \ldots, v_6, w_1, \ldots, w_6\}$$

and

$$\begin{aligned} E(G') = \quad & \{u_iv_j |\ 1 \leq i \leq 3, 1 \leq j \leq 6\} \cup \{w_1w_2, w_3w_4\} \cup \{w_iw_j |\ 1 \leq i \leq 4, j = 5, 6\} \\ & \cup \{v_1w_3, v_1w_4, v_2w_1, v_2w_2, v_3w_5, v_3w_6, v_4w_3, v_4w_4, v_5w_1, v_5w_2, v_6w_5, v_6w_6\}, \end{aligned}$$

by the Euler characteristic formula, there are 25 faces when drawing G' on a torus. Fix a representation of G' and let $\{F'_1, \ldots, F'_{25}\}$ be the set of faces of G' corresponding to that representation. Since $G'' \simeq K_{3,6}$, this graph has 9 faces (see Remark 1.4). Let $F''_1, \ldots, F''_9$ be the faces of G'' obtained by deleting $w_1, \ldots, w_6$ and all edges incident with $w_1, \ldots, w_6$ from this representation. Again, $\{F'_1, \ldots, F'_{25}\}$ can be recovered by inserting $w_1, \ldots, w_6$ and all edges incident with $w_1, \ldots, w_6$ into the representation corresponding to $\{F''_1, \ldots, F''_9\}$. We note that $w_iw_j \in E(G')$ for $i = 1, \ldots, 4$ and $j = 5, 6$. Therefore all w_i should be inserted into the same face, say F''_m, of G'' to avoid crossings. Moreover, since $v_1w_3, v_2w_4, v_3w_5, v_4w_4, v_5w_2, v_6w_5 \in E(G')$, it follows that $v_1, v_2, \ldots, v_6$ are all vertices of F''_m. This contradicts the fact that the boundary of F''_m is a 4-cycle. Thus, we conclude that $\gamma(G) \geq 2$.

Case 2 : $y^2 = 0$. In this case, we may further assume that $z^2 = 0$. Since $z\mathfrak{m} \neq \{0\}$ by assumption, we have $yz \neq 0$. Consequently, $x^2 = yz$ and $xy = xz = y^2 = z^2 = 0$. Let $u_1 = x^2$, $u_2 = x$, $u_3 = x + x^2$, $v_1 = y$, $v_2 = z$, $v_3 = y + z$, $v_4 = y + x^2$, $v_5 = z + x^2$, $v_6 = y + z + x^2$, $w_1 = x + y$, $w_2 = x + y + x^2$, $w_3 = x + z$, $w_4 = x + z + x^2$, $w_5 = x + y + z$ and $w_6 = x + y + z + x^2$. Then $u_i \cdot v_j = 0$ for every i, j, so that $K_{3,6} \subseteq \Gamma(R)$. Therefore $\gamma(\Gamma(R)) \geq \gamma(K_{3,6}) = 1$. Let $G = \Gamma(R)$, $G' = G - \{w_5, w_6\} - \{u_1w_1, u_1w_2, u_1w_3, u_1w_4, u_1u_2, u_1u_3, v_1v_4, v_2v_5, v_3v_6\}$, and $G'' = G' - \{w_1, w_2, w_3, w_4\}$. Thus $G'' \simeq K_{3,6}$.

Suppose that $\gamma(G) = 1$, and so $\gamma(G') = 1$. Since

$$V(G') = \{u_1, u_2, u_3, v_1, v_2, \ldots, v_6, w_1, \ldots, w_4\}$$

and

$$\begin{aligned} E(G') = \quad & \{u_i v_j \mid 1 \le i \le 3, 1 \le j \le 6\} \cup \{w_1w_3, w_1w_4, w_2w_3, w_2w_4\} \\ & \cup \{v_1w_1, v_1w_2, v_2w_3, v_2w_4, v_4w_1, v_4w_2, v_5w_3, v_5w_4\}, \end{aligned}$$

G' has 17 faces. By a similar deletion and insertion argument as before, fix a representation of G' and let $\{F'_1, \ldots, F'_{17}\}$ be the set of faces of G' corresponding to this representation. Let $\{F''_1, \ldots, F''_9\}$ be the set of faces of G'' obtained by deleting $w_1, \ldots, w_4$ and all edges incident with $w_1, \ldots, w_4$ from G'. Since $w_1w_3, w_2w_3, w_1w_4, w_2w_4 \in E(G')$, w_1, w_2, w_3, and w_4 should be inserted into the same face, say F''_n, of G'' in the recovering process from G'' to G' to avoid crossings. We note that $v_1w_1, v_2w_3, v_4w_1, v_5w_4 \in E(G')$ and therefore v_1, v_2, v_4, v_5 are the four vertices of F''_n. Denote $e_1 = v_1w_1$,$e_2 = v_1w_2$, $e_3 = v_2w_3$, $e_4 = v_2w_4$, $e_5 = w_1w_3$, $e_6 = w_1w_4$, $e_7 = w_2w_3$, and $e_8 = w_2w_4$. Then we obtain Figure 2 by inserting $w_1, \ldots, w_4$ and $e_1, \ldots, e_4$ into F''_n. However, from Figure 2 we see that there is no way to insert $e_5, \ldots, e_8$ into F''_n without crossings, a contradiction. Therefore, we conclude that $\gamma(G) \ge 2$. □



Figure 2

Proposition 2.3. *Let $(R, \mathfrak{m})$ be a local ring. If $|R| = 16$, $|R/\mathfrak{m}| = 2$ and $\mathfrak{m}^3 = 0$, then $\gamma(\Gamma(\mathbb{Z}_2 \times R)) \ge 2$.*

Proof. By hypothesis, we have $|\mathfrak{m}| = 8$. If $|\mathfrak{m}^2| = 4$, then $\mathfrak{m}$ is principal, and so is $\mathfrak{m}^2$. This implies that $|\mathfrak{m}^3| = 2$, a contradiction. So $\mathfrak{m}^2 = 0$ or $|\mathfrak{m}^2| = 2$. If $\mathfrak{m}^2 = 0$, then the seven non-zero elements of $\mathfrak{m}$ are all zero-divisors of R. Write $\mathfrak{m} - \{0\} = \{a_1, \ldots, a_7\}$ and let $u_i = (0, a_i)$ and $v_i = (1, a_i)$ for $i = 1, \ldots, 7$. Then $u_i \cdot v_j = 0$ for every $1 \le i, j \le 7$, so that $K_{7,7} \subseteq \Gamma(\mathbb{Z}_2 \times R)$, and it follows that $\gamma(\Gamma(\mathbb{Z}_2 \times R)) \ge \gamma(K_{7,7}) = 7$. Next, we assume that $|\mathfrak{m}^2| = 2$, so that $dim_{R/\mathfrak{m}}\mathfrak{m}/\mathfrak{m}^2 = 2$ and $dim_{R/\mathfrak{m}}\mathfrak{m}^2/\mathfrak{m}^3 = 1$. Let $\mathfrak{m} = (x, y)$ for some $x, y \in \mathfrak{m} - \mathfrak{m}^2$. We now consider two cases.

Case 1. Suppose $x^2 = y^2 = 0$. In this case, $xy \ne 0$ as $\mathfrak{m}^2 \ne 0$. Let $u_1 = (0, xy)$, $u_2 = (1, xy)$, $u_3 = (1, 0)$, $v_1 = (0, x)$, $v_2 = (0, x + xy)$, $v_3 = (0, y)$, $v_4 = (0, y + xy)$, $v_5 = (0, x + y)$, $v_6 = (0, x + y + xy)$, $w_1 = (1, x)$, $w_2 = (1, x + xy)$, $w_3 = (1, y)$, $w_4 = (1, y + xy)$, $w_5 = (1, x + y)$ and $w_6 = (1, x + y + xy)$; then $u_i \cdot v_j = 0$ for every i, j, so that $K_{3,6} \subseteq \Gamma(\mathbb{Z}_2 \times R)$, it follows that $\gamma(\Gamma(\mathbb{Z}_2 \times R)) \ge \gamma(K_{3,6}) = 1$. Write $G = \Gamma(\mathbb{Z}_2 \times R)$. Note that if $z \notin \mathfrak{m}$ then $(0, z) \in V(G)$ and $deg\,((0, z)) = 1$. So $V(\widetilde{G})$ consists of all u_i, v_j and w_k as defined above. Let $G' = \widetilde{G} - \{w_3, w_4, w_5, w_6\} - \{u_1u_2, u_1u_3, v_1v_2, v_3v_4, v_5v_6\}$ and $G'' = G' - \{w_1, w_2\}$, so that $G'' \simeq K_{3,6}$.

Suppose that $\gamma(G) = 1$. Since $\gamma(G'') \le \gamma(G') \le \gamma(G)$, we have $\gamma(G') = 1$. Since

$$V(G') = \{u_1, u_2, u_3, v_1, \ldots, v_6, w_1, w_2\}$$

and

$$E(G') = \{u_iv_j \mid 1 \le i \le 3, 1 \le j \le 6\} \cup \{v_iw_j \mid i,j = 1,2\} \cup \{u_1w_1, u_1w_2\},$$

G' has 13 faces. Fix a representation of G' and let $\{F'_1, \ldots, F'_{13}\}$ be the set of faces of G' corresponding to this representation. Let $\{F''_1, \ldots, F''_9\}$ be the set of faces of G'' obtained by deleting w_1, w_2 and all edges incident with w_1, w_2 from G'. Therefore there are faces F''_{t_1}, F''_{t_2} so that inserting w_1, w_2 and all edges incident with w_1, w_2 into these faces, we are able to recover the set of faces $\{F'_1, \ldots, F'_{13}\}$. Since $w_1v_1 = w_2v_1 = w_1v_2 = w_2v_2 \in V(G')$, w_1 and w_2 should be inserted into the same face, say F''_l, of G''. Let $e_1 = v_1w_1$, $e_2 = v_1w_2$, $e_3 = v_2w_1$, $e_4 = v_2w_2$, $e_5 = u_1w_1$ and $e_6 = u_1w_2$. After inserting w_1, w_2 and e_1, e_2, e_3, e_4 into F'_l we obtain Figure 3 as below. From the figure, we see that there is no way to insert e_5, e_6 into F'_l without any crossings. Thus, we conclude that $\gamma(G) \ge 2$.



Figure 3

Case 2. Suppose $x^2 \neq 0$. In this case, we may assume $xy = 0$. Otherwise, $x^2 = xy$, and after replacing y by $y - x$ we have $xy = 0$. We note that either $y^2 = 0$ or $y^2 = x^2$ as $|\mathfrak{m}^2| = 2$. Assume that $y^2 = 0$. Let $u_1 = (0, y)$, $u_2 = (0, x^2)$, $u_3 = (0, y + x^2)$, $v_1 = (0, x + y)$, $v_2 = (1, x + y)$, $v_3 = (0, x)$, $v_4 = (1, x)$, $v_5 = (0, x + x^2)$, $v_6 = (0, x + y + x^2)$ and $v_7 = (1, x + x^2)$; then $u_i \cdot v_j = 0$ for every i, j, so that $K_{3,7} \subseteq \Gamma(\mathbb{Z}_2 \times R)$, and it follows that $\gamma(\Gamma(\mathbb{Z}_2 \times R)) \ge \gamma(K_{3,7}) = 2$. Therefore, it remains to discuss the case when $y^2 = x^2$. Let $u_1 = (0, x^2)$, $u_2 = (1, x^2)$, $u_3 = (1, 0)$, $v_1 = (0, y)$, $v_2 = (0, y + x^2)$, $v_3 = (0, x)$, $v_4 = (0, x + x^2)$, $v_5 = (0, x + y)$, $v_6 = (0, x + y + x^2)$, $w_1 = (1, x)$, $w_2 = (1, x + x^2)$, $w_3 = (1, y)$, $w_4 = (1, y + x^2)$, $w_5 = (1, x + y)$ and $w_6 = (1, x + y + x^2)$. Then $u_i \cdot v_j = 0$ for every i, j, so that $K_{3,6} \subseteq \Gamma(\mathbb{Z}_2 \times R)$, and it follows that $\gamma(\Gamma(\mathbb{Z}_2 \times R)) \ge \gamma(K_{3,6}) = 1$.Let $G = \Gamma(\mathbb{Z}_2 \times R)$ and $G^* = \widetilde{G} - \{w_3, w_4, w_5, w_6\} - \{u_1u_2, u_1u_3, v_1v_3, v_1v_4, v_2v_3, v_2v_4, v_5v_6\}$. We observe that G^* is isomorphic to the graph G' obtained in Case 1. Therefore $\gamma(G^*) \ge 2$ and thus $\gamma(G) \ge 2$. □

Proposition 2.4. *Let $(R, \mathfrak{m})$ be a local ring such that $|R| = p^n$, where p is prime and $n \in \mathbb{N}$. If $\mathfrak{m}^{n-1} \neq 0$, then $\mathfrak{m}$ is principal and $\Gamma(R) \simeq \Gamma(\mathbb{Z}_{p^n})$. Moreover, for any ring S, one has that $\Gamma(S \times R) \simeq \Gamma(S \times \mathbb{Z}_{p^n})$.*

Proof. We note that $\mathfrak{m}^{k+1} \subsetneq \mathfrak{m}^k$ if $\mathfrak{m}^k \neq 0$, so that $|\mathfrak{m}^k| \le p^{n-k}$ for $k \le n$. Since $\mathfrak{m}^{n-1} \neq 0$ we have $|\mathfrak{m}^k| = p^{n-k}$, so that $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 1$ and therefore $\mathfrak{m}$ is principal. Suppose $\mathfrak{m} = (x)$ and let $\widetilde{\mathfrak{m}}_k = \mathfrak{m}^k - m^{k+1}$. Then $\widetilde{\mathfrak{m}}_k = \{c_1x^k, \ldots, c_lx^k\}$, where $l = p^{n-k-1}(p-1)$ and each c_i is a unit in R. Observe that for $u_i \in \widetilde{\mathfrak{m}}_i$ and $v_j \in \widetilde{\mathfrak{m}}_j$, $u_i \cdot v_j = 0$ if and only if $i + j \ge n$. Thus, $\Gamma(R)$ is uniquely determined. In particular, $\Gamma(R) \simeq \Gamma(\mathbb{Z}_{p^n})$ as $\mathbb{Z}_{p^n}$ satisfies these assumptions. Moreover, let $\psi : R \to \mathbb{Z}_{p^n}$ be a bijective map such that $\psi(\widetilde{\mathfrak{m}}_k) = (\widetilde{\mathfrak{n}}_k)$ for each $k < n$, where $\mathfrak{n} = (p)$ is the maximal ideal of $\mathbb{Z}_{p^n}$. It is easy to see that $\phi : S \times R \to S \times \mathbb{Z}_{p^n}$ with $\phi(a, b) = (a, \psi(b))$ induces an isomorphism between $\Gamma(S \times R)$ and $\Gamma(S \times \mathbb{Z}_{p^n})$. □

We now briefly turn our attention to the case where $\Gamma(R)$ is planar. In the following three examples, we show by explicit representations that the zero-divisor graphs of the listed local rings are planar.

Example 2.5. (a) $\Gamma(\mathbb{Z}_{27})$ is isomorphic to G_1 as in Figure 4-1. (b) $\Gamma(\mathbb{Z}_{16})$ is isomorphic to G_2 as shown in Figure 4-2.

Proof. (a) In $\mathbb{Z}_{27}$, let $u_1 = \bar{3}$, $u_2 = \bar{6}$, $u_3 = \bar{12}$, $u_4 = \bar{15}$, $u_5 = \bar{21}$, $u_6 = \bar{24}$, $v_1 = \bar{9}$, $v_2 = \bar{18}$. Since $u_i \cdot v_j = 0$ and $v_1 \cdot v_2 = 0$, $\Gamma(\mathbb{Z}_{27})$ is isomorphic to G_1 as shown in Figure 4-1.

(b) In $\mathbb{Z}_{16}$, let $u = \bar{8}$, $v_1 = \bar{2}$, $v_2 = \bar{6}$, $v_3 = \bar{4}$, $v_4 = \bar{12}$, $v_5 = \bar{10}$, $v_6 = \bar{14}$. Since $u \cdot v_j = 0$ and $v_3 \cdot v_4 = 0$, $\Gamma(\mathbb{Z}_{16})$ is isomorphic to G_2 as shown in Figure 4-2. □



Figure 4-1



Figure 4-2

Example 2.6. Let R be one of the following local rings: (a) $\mathbb{Z}_2[x,y]/(x^3, xy, y^2 - x^2)$, (b) $\mathbb{Z}_4[x]/(x^3, x^2 - 2x)$, (c) $\mathbb{Z}_4[x,y]/(x^3, x^2 - 2, xy, y^2 - 2)$, or (d) $\mathbb{Z}_8[x]/(x^2 - 4, 2x)$. Then $\Gamma(R)$ is planar and is isomorphic to G_3 as shown in Figure 5-1.

Proof. We briefly sketch the details for each case.

(a) If $R = \mathbb{Z}_2[x,y]/(x^3, xy, y^2 - x^2)$, then $Z(R)^* = \{\bar{x}, \bar{y}, \bar{x}^2, \bar{x} + \bar{x}^2, \bar{y} + \bar{x}^2, \bar{x} + \bar{y}, \bar{x} + \bar{y} + \bar{x}^2\}$. Let $u = \bar{x}^2$, $v_1 = \bar{x}$, $v_2 = \bar{y}$, $v_3 = \bar{x} + \bar{x}^2$, $v_4 = \bar{y} + \bar{x}^2$, $v_5 = \bar{x} + \bar{y}$, and $v_6 = \bar{x} + \bar{y} + \bar{x}^2$. Then $\Gamma(R) \simeq G_3$ as shown in Figure 5-1.

(b) If $R = \mathbb{Z}_4[x]/(x^3, x^2 - 2x)$, then $Z(R)^* = \{\bar{2}, \bar{x}, \bar{2x}, \bar{2} + \bar{2x}, \bar{3x}, \bar{2} + \bar{x}, \bar{2} + \bar{3x}\}$. Let $u = \bar{2x}$, $v_1 = \bar{x}$, $v_2 = \bar{2} + \bar{x}$, $v_3 = \bar{3x}$, $v_4 = \bar{2} + \bar{3x}$, $v_5 = \bar{2}$ and $v_6 = \bar{2} + \bar{2x}$. It then follows that $\Gamma(R) \simeq G_3$ as shown in Figure 5-1.

(c) If $R = \mathbb{Z}_4[x,y]/(x^3, x^2 - 2, xy, y^2 - 2)$, then $Z(R)^* = \{\bar{2}, \bar{x}, \bar{y}, \bar{x} + \bar{2}, \bar{y} + \bar{2}, \bar{x} + \bar{y}, \bar{x} + \bar{y} + \bar{2}\}$. Let $u = \bar{2}$, $v_1 = \bar{x}$, $v_2 = \bar{y}$, $v_3 = \bar{x} + \bar{2}$, $v_4 = \bar{y} + \bar{2}$, $v_5 = \bar{x} + \bar{y}$ and $v_6 = \bar{x} + \bar{y} + \bar{2}$. Then $\Gamma(R) \simeq G_3$ as shown in Figure 5-1.

(d) If $R = \mathbb{Z}_8[x]/(x^2 - 4, 2x)$, then $Z(R)^* = \{\bar{2}, \bar{4}, \bar{6}, \bar{x}, \bar{2} + \bar{x}, \bar{4} + \bar{x}, \bar{6} + \bar{x}\}$. Let $u = \bar{4}$, $v_1 = \bar{2}$, $v_2 = \bar{x}$, $v_3 = \bar{6}$, $v_4 = \bar{4} + \bar{x}$, $v_5 = \bar{2} + \bar{x}$ and $v_6 = \bar{6} + \bar{x}$. Then $\Gamma(R) \simeq G_3$ as shown in Figure 5-1. □



Figure 5-1



Figure 5-2

Example 2.7. If R is one of the following local rings: (a) $\mathbb{Z}_4[x,y]/(x^2, y^2, xy - 2)$, (b) $\mathbb{Z}_2[x,y]/(x^2, y^2)$, or (c) $\mathbb{Z}_4[x]/(x^2)$, then $\Gamma(R)$ is planar and is isomorphic to G_4 as shown in Figure 5-2.

Proof. (sketch)

(a) If $R = \mathbb{Z}_4[x,y]/(x^2, y^2, xy-2)$, then $Z(R)^* = \{\bar{2}, \bar{x}, \bar{y}, \bar{x}+\bar{2}, \bar{y}+\bar{2}, \bar{x}+\bar{y}, \bar{x}+\bar{y}+\bar{2}\}$. Let $u = \bar{2}$, $v_1 = \bar{x}$, $v_2 = \bar{x}+\bar{2}$, $v_3 = \bar{y}$, $v_4 = \bar{y}+\bar{2}$, $v_5 = \bar{x}+\bar{y}$ and $v_6 = \bar{x}+\bar{y}+\bar{2}$. Then $\Gamma(R) \simeq G_4$ as shown in Figure 5-2.

(b) If $R = \mathbb{Z}_2[x,y]/(x^2, y^2)$, then $Z(R)^* = \{\bar{x}, \bar{y}, \bar{x}\bar{y}, \bar{x}+\bar{x}\bar{y}, \bar{y}+\bar{x}\bar{y}, \bar{x}+\bar{y}, \bar{x}+\bar{y}+\bar{x}\bar{y}\}$. Let $u = \bar{x}\bar{y}$, $v_1 = \bar{x}$, $v_2 = \bar{x}+\bar{x}\bar{y}$, $v_3 = \bar{y}$, $v_4 = \bar{y}+\bar{x}\bar{y}$, $v_5 = \bar{x}+\bar{y}$ and $v_6 = \bar{x}+\bar{y}+\bar{x}\bar{y}$. Thus $\Gamma(R) \simeq G_4$ as shown in Figure 5-2.

(c) If $R = \mathbb{Z}_4[x]/(x^2)$, then $Z(R)^* = \{\bar{2}, \bar{x}, 2\bar{x}, \bar{2}+2\bar{x}, 3\bar{x}, \bar{2}+\bar{x}, \bar{2}+3\bar{x}\}$. Let $u = 2\bar{x}$, $v_1 = \bar{2}$, $v_2 = \bar{2}+2\bar{x}$, $v_3 = \bar{x}$, $v_4 = 3\bar{x}$, $v_5 = \bar{2}+\bar{x}$ and $v_6 = \bar{2}+3\bar{x}$. Then $\Gamma(R) \simeq G_4$ as shown in Figure 5-2. □

From Lemma 1.1, we see that complete graphs on 7 vertices are interesting in the genus one setting. Thus, we discuss some examples of finite local rings whose zero-divisor graphs have seven vertices.

Remark 2.8. Suppose G is a simple graph such that $|V(G)| = 7$ and G contains a subgraph isomorphic to $K_{3,4}$, i.e., $K_{3,4} \subseteq G \subseteq K_7$. Then by Lemma 1.1, 1.2 and Remark 1.3, we have $1 = \gamma(K_{3,4}) \leq \gamma(G) \leq \gamma(K_7) = 1$. In particular, for the 4-partite graph $K_{1,1,1,4}$ one has $\gamma(K_{1,1,1,4}) = 1$.

(2) Consider $G_5 = \Gamma(\mathbb{Z}_{32})$. Then $V(G_5) = \{\bar{2}, \bar{4}, \bar{6}, \ldots, \bar{30}\}$. Let $u_1 = \bar{8}, u_2 = \bar{16}, u_3 = \bar{24}$, $v_1 = \bar{4}$, $v_2 = \bar{12}$, $v_3 = \bar{20}$, $v_4 = \bar{28}$, $w_1 = \bar{2}$, $w_2 = \bar{6}$, $w_3 = \bar{10}$, $w_4 = \bar{14}$, $w_5 = \bar{18}$, $w_6 = \bar{22}$, $w_7 = \bar{26}$, and $w_8 = \bar{30}$. We note that $deg(w_i) = 1$ for each i, so that $\widetilde{G_5} = G_5 - \{w_1, \ldots, w_8\}$. Moreover, since $u_i \cdot u_j = 0$, $u_i \cdot v_j = 0$ and $v_i \cdot v_j \neq 0$ for all pairs i, j, we have $\widetilde{G_5} \simeq K_{1,1,1,4}$. Therefore, $\gamma(G_5) = \gamma(\widetilde{G_5}) = 1$ and thus G_5 is isomorphic to $K_{1,1,1,4}$ with 8 single pendant edges.[†]

Example 2.9. Let R be one of the following local rings: (a) $\mathbb{Z}_2[x,y]/(x^3, xy, y^2)$, (b) $\mathbb{Z}_4[x]/(x^3, 2x)$, (c) $\mathbb{Z}_4[x,y]/(x^3, x^2-2, xy, y^2)$, or(d) $\mathbb{Z}_8[x]/(x^2, 2x)$. Then $\Gamma(R) \simeq K_{1,1,1,4}$ and $\gamma(\Gamma(R)) = 1$.

Proof. (a) If $R = \mathbb{Z}_2[x,y]/(x^3, xy, y^2)$, then $Z(R)^* = \{\bar{x}, \bar{y}, \bar{x}^2, \bar{x}+\bar{x}^2, \bar{y}+\bar{x}^2, \bar{x}+\bar{y}, \bar{x}+\bar{y}+\bar{x}^2\}$. Let $u_1 = \bar{y}$, $u_2 = \bar{x}^2$, $u_3 = \bar{y}+\bar{x}^2$, $v_1 = \bar{x}$, $v_2 = \bar{x}+\bar{x}^2$, $v_3 = \bar{x}+\bar{y}$ and $v_4 = \bar{x}+\bar{y}+\bar{x}^2$. It is easy to see that for all i, j we have $u_i \cdot u_j = 0$, $u_i \cdot v_j = 0$, and $v_i \cdot v_j \neq 0$. Thus $\Gamma(R) \simeq K_{1,1,1,4}$ and $\gamma(\Gamma(R)) = 1$ by Remark 2.8 (1).

(b) If $R = \mathbb{Z}_4[x]/(x^3, 2x)$, then $Z(R)^* = \{2, \bar{x}, 2+\bar{x}, \bar{x}^2, 2+\bar{x}^2, \bar{x}+\bar{x}^2, 2+\bar{x}+\bar{x}^2\}$. Let $u_1 = 2$, $u_2 = \bar{x}^2$, $u_3 = 2+\bar{x}^2$, $v_1 = \bar{x}$, $v_2 = 2+\bar{x}$, $v_3 = \bar{x}+\bar{x}^2$ and $v_4 = 2+\bar{x}+\bar{x}^2$. As in the previous case, it is clear that $\Gamma(R) \simeq K_{1,1,1,4}$.

(c) If $R = \mathbb{Z}_4[x,y]/(x^3, x^2-2, xy, y^2)$, then $Z(R)^* = \{2, \bar{x}, \bar{y}, \bar{x}+2, \bar{y}+2, \bar{x}+\bar{y}, \bar{x}+\bar{y}+2\}$. Let $u_1 = 2$, $u_2 = \bar{y}$, $u_3 = \bar{y}+2$, $v_1 = \bar{x}$, $v_2 = \bar{x}+2$, $v_3 = \bar{x}+\bar{y}$ and $v_4 = \bar{x}+\bar{y}+2$. Again, it follows immediately that $\Gamma(R) \simeq K_{1,1,1,4}$.

(d) If $R = \mathbb{Z}_8[x]/(x^2, 2x)$, then $Z(R)^* = \{2, 4, 6, \bar{x}, 2+\bar{x}, 4+\bar{x}, 6+\bar{x}\}$. Let $u_1 = 4$, $u_2 = \bar{x}$, $u_3 = 4+\bar{x}$, $v_1 = 2$, $v_2 = \bar{x}+2$, $v_3 = 6$ and $v_4 = \bar{x}+6$. Once again, it is easy to see that $\Gamma(R) \simeq K_{1,1,1,4}$ and $\gamma(\Gamma(R)) = 1$ by Remark 2.8. □

We end this section by discussing how the genus number of zero-divisor graphs behaves with respect to products.

Example 2.10. Let $R = \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{F}_4$, and denote the elements of $\mathbb{F}_4$ as x_0, x_1, x_2, and x_3. We note that the reduction of $\Gamma(R)$, denoted by G_6, consists of 11 vertices, namely $u_1 = (0,0,x_1), u_2 = (0,0,x_2), u_3 = (0,0,x_3), v_1 = (1,0,x_0), v_2 = (1,1,x_0), v_3 = (1,2,x_0), v_4 = (0,1,x_0), v_5 = (0,2,x_0), w_1 = (1,0,x_1), w_2 = (1,0,x_2), w_3 = (1,0,x_3)$, with edge set $E(G_6) = \{\, u_i v_j \mid i = 1,2,3; j = 1,2,3,4,5 \,\} \cup \{\, v_r v_j \mid r = 4,5; j = 1,2,3 \,\} \cup \{\, v_r w_j \mid r = 4,5; j = 1,2,3 \,\}$. Since $K_{3,5} \subseteq G_6$, we have that $\gamma(G_6) \geq 1$. The following embedding explicitly shows that $\gamma(G_6) = 1$.

[†] For the representation of the 4-partite graph $K_{1,1,1,4}$, refer to Figure 5(b) in [20].



Figure 6

Proposition 2.11. *Let $(R, \mathfrak{m})$ be a finite local ring such that $|\mathfrak{m}| = 4$ and $\mathfrak{m}^2 = 0$. Then $\gamma(\Gamma(\mathbb{Z}_2 \times R)) = 1$.*

Proof. Let $G = \Gamma(\mathbb{Z}_2 \times R)$ and let a_1, a_2, a_3 be distinct nonzero elements in $\mathfrak{m}$. Let $u_i = (0, a_i)$ and $v_i = (1, a_i)$ for $i = 1, 2, 3$. Then, $u_i \cdot v_j = 0$ for each pair i, j, so that $K_{3,3} \subseteq G$ and therefore $\gamma(G) \geq \gamma(K_{3,3}) = 1$. On the other hand, let $w_1 = (1, a_1)$, $w_2 = (0, a_1)$, $w_3 = (1, a_2)$, $w_4 = (0, a_2)$, $w_5 = (1, a_3)$, $w_6 = (0, a_3)$, and $w_7 = (1, 0)$. Therefore $V(\widetilde{G}) = \{w_1, \ldots, w_7\}$. Let $\phi : \widetilde{G} \to \Gamma(\mathbb{Z}_{32})$ be the map obtained by sending w_i to $4i$ for every i. It is easy to see that $\widetilde{G}$ is a subgraph of $\Gamma(\mathbb{Z}_{32})$ via ϕ, so that $\gamma(G) = \gamma(\widetilde{G}) \leq \gamma(\Gamma(\mathbb{Z}_{32})) = 1$ by [20, Theorem 4.5]. Thus, we conclude $\gamma(G) = 1$. □

Remark 2.12. Let $\mathbb{F}_q$ denote the finite field with q elements and let $\psi : \mathbb{F}_q \to \mathbb{Z}_q$ be any bijective map such that $\psi(0) = 0$. Let R be a ring and define a map $\phi : R \times \mathbb{F}_q \to R \times \mathbb{Z}_q$ such that $\phi((a, b)) = (a, \psi(b))$. Then ϕ induces an embedding from $\Gamma(R \times \mathbb{F}_q)$ to $\Gamma(R \times \mathbb{Z}_q)$. Therefore we conclude that $\gamma(\Gamma(R \times \mathbb{F}_q)) \leq \gamma(\Gamma(R \times \mathbb{Z}_q))$.

3 Toroidal zero-divisor graphs

The main goal of this section is to determine all finite rings R whose zero-divisor graphs are of genera at most one. To do this we examine the characteristic of R, denoted by $char(R)$, the cardinality of the residue field $|R/\mathfrak{m}|$ (when R is local), and the number of irreducible components of R. We begin this section with a few results related to the characteristic of a ring R.

Lemma 3.1. *Let $(R, \mathfrak{m})$ be a finite local ring such that $|R| = q^2$ and $|R/\mathfrak{m}| = q = p^d$ for some prime p and $d \in \mathbb{N}$. If $char(R) = p^2$, then $R \simeq \mathbb{Z}_{p^2}[x]/(f(x))$, where f is a monic polynomial of $\mathbb{Z}_{p^2}[x]$ of degree d such that the image of f in $\mathbb{Z}_p[x]$ is irreducible.*

Proof. By hypothesis, we have $|\mathfrak{m}| = q$. Since $\mathfrak{m}/\mathfrak{m}^2$ is a vector space over $R/\mathfrak{m}$, it follows that $|\mathfrak{m}/\mathfrak{m}^2| = q^r$, where $r = \dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2$. Therefore $\mathfrak{m}^2 = 0$ and $r = 1$, and this implies that $\mathfrak{m} = (x)$ for every $x \in \mathfrak{m} - \{0\}$. Moreover, since $char(R) = p^2$ and $\mathbb{Z}_{p^2} \subset R$, we have that R is a finitely generated $\mathbb{Z}_{p^2}$-algebra. That is, $R = \mathbb{Z}_{p^2}[u_1, \ldots, u_n]$ for some $u_i \in R$. We note that $\mathfrak{m} = (p)$ so that $\mathbb{F}_q \simeq R/\mathfrak{m} = \mathbb{Z}_{p^2}[u_1, \ldots, u_n]/(p) = \mathbb{Z}_p[\bar{u}_1, \ldots, \bar{u}_n]$, where $\bar{u}_i$ is the image of u_i in $\mathbb{Z}_{p^2}[u_1, \ldots, u_n]/(p)$. Since $\mathbb{F}_q$ is a simple extension of $\mathbb{Z}_p$, there exists $u \in R$ such that $\mathbb{Z}_p[\bar{u}_1, \ldots, \bar{u}_n] = \mathbb{Z}_p[\bar{u}]$. Hence for any $w \in R$, there exists $g_w(x) \in \mathbb{Z}_{p^2}[x]$ such that

$$w \equiv g_w(u) \pmod{\mathfrak{m}}.$$

This implies that there exists $g_w^*(x_1, \ldots, x_n) \in \mathbb{Z}_{p^2}[x_1, \ldots, x_n]$ such that

$$w - g_w(u) - pg_w^*(u_1, \ldots, u_n) = 0. \tag{1}$$

Since $char(R) = p^2$, it follows from (1) that

$$pw = pg_w(u). \tag{2}$$

That is, for every $w \in R$, there exists $g_w(x) \in \mathbb{Z}_{p^2}[x]$ such that $pw = pg_w(u)$. Replacing w by $g_w^*(u_1, \ldots, u_n)$, we see from (2) that there exists $g_w^{**}(x) \in \mathbb{Z}_{p^2}[x]$, such that

$$pg_w^*(u_1, \ldots, u_n) = pg_w^{**}(u). \tag{3}$$

It then follows from (1) and (3) that

$$w = g_w(u) + pg_w^{**}(u) \in \mathbb{Z}_{p^2}[u],$$

so that $R = \mathbb{Z}_{p^2}[u]$. Next, we proceed to show that there exists a monic, d-degree polynomial $f(x) \in \mathbb{Z}_{p^2}[x]$ such that $f(u) = 0$.

Since $[\,\mathbb{Z}_p[\bar{u}] : \mathbb{Z}_p] = d$, there exists a monic, irreducible polynomial $h(x) \in \mathbb{Z}_p[x]$ of degree d such that $h(\bar{u}) = 0$. This implies that there exists a monic polynomial $l(x) \in \mathbb{Z}_{p^2}[x]$ of degree d such that $\overline{l(u)} = h(\bar{u}) = 0$, i.e., $l(u) \in \mathfrak{m}$, so that $l(u) = pl_1(u)$ for some $l_1(x) \in \mathbb{Z}_{p^2}[x]$. If $\overline{l_1(u)} = 0$ then $l_1(u) = pl_2(u)$ for some $l_2(x) \in \mathbb{Z}_{p^2}[x]$. This implies that $l(u) = p^2 l_2(u) = 0$. Set $f(x) = l(x)$ and then we are done. Otherwise, since $[\,\mathbb{Z}_p[\bar{u}] : \mathbb{Z}_p] = d$, $\overline{l_1(u)} = h_2(\bar{u})$ for some $h_2(x) \in \mathbb{Z}_p[x]$ with $\deg h_2(x) < d$. Similarly, there exists $l_2(x) \in \mathbb{Z}_{p^2}[x]$ such that $\deg l_2(x) = \deg h_2(x) < d$ and $\overline{l_2(u)} = h_2(\bar{u})$. Therefore $\overline{l_1(u)} = \overline{l_2(u)}$ and this implies that $l_1(u) - l_2(u) \in \mathfrak{m}$, so that $pl_1(u) = pl_2(u)$. Let $f(x) = l(x) - pl_2(x) \in \mathbb{Z}_{p^2}[x]$. Then $f(u) = l(u) - pl_2(u) = l(u) - pl_1(u) = 0$, and $\deg f(x) = d$ and the image of f in $\mathbb{Z}_p[x]$ (which is equal to $h(x)$) is irreducible.

Let $\phi : \mathbb{Z}_{p^2}[x] \to \mathbb{Z}_{p^2}[u]$ be the natural ring homomorphism which maps x to u. We note that ϕ is surjective and that $f(x) \in \ker \phi$, so we have

$$q^2 = |\,\mathbb{Z}_{p^2}[u]\,| = |\,\mathbb{Z}_{p^2}[x]/\ker \phi\,| \leq |\,\mathbb{Z}_{p^2}[x]/(f(x))\,|.$$

However, since $\mathbb{Z}_{p^2}[x]/(f(x))$ is a free $\mathbb{Z}_{p^2}$-module of rank d, it has $(p^2)^d = q^2$ elements. Thus $\ker \phi = (f(x))$ so that $R = \mathbb{Z}_{p^2}[u] \simeq \mathbb{Z}_{p^2}[x]/(f(x))$ which completes the proof. □

Applying Lemma 3.1 for the cases $(q, p, d) = (8, 2, 3)$ and $(q, p, d) = (4, 2, 2)$ we obtain the following two corollaries.

Corollary 3.2. *Let $(R, \mathfrak{m})$ be a finite local ring such that $|R| = 64$ and $|R/\mathfrak{m}| = 8$. If $char(R) = 4$, then $R \simeq \mathbb{Z}_4[x]/(x^3 + x + 1)$.*

Proof. We note that there are two irreducible polynomials in $\mathbb{Z}_2[x]$ of degree 3, namely $x^3 + x^2 + 1$ and $x^3 + x + 1$. Applying Lemma 3.1 for $(q, p, d) = (4, 2, 2)$, we see that $R \simeq \mathbb{Z}_4[x]/(f(x))$, where $\overline{f(x)} = x^3+x+1$ or $\overline{f(x)} = x^3+x^2+1$. Let $g_1(x) = x^3+x+1$, $g_2(x) = x^3+x-1$, $g_3(x) = x^3+2x^2+x-1$, $g_4(x) = x^3+2x^2+x+1$, $g_5(x) = x^3+x^2-1$, $g_6(x) = x^3+x^2+1$, $g_7(x) = x^3-x^2-1$, $g_8(x) = x^3+x^2+1$, $h_1(x) = x^3-x+1$, $h_2(x) = x^3-x-1$, $h_3(x) = x^3+2x^2-x-1$, $h_4(x) = x^3+2x^2-x+1$, $h_5(x) = x^3-x^2+2x+1$, $h_6(x) = x^3-x^2+2x-1$, $h_7(x) = x^3+x^2+2x+1$, and $h_8(x) = x^3+x^2+2x-1 \in \mathbb{Z}_4[x]$. Then $f(x)$ is either $g_i[x]$ or $h_j[x]$. We note that $\mathbb{Z}_4[x]/(g_i[x])$ are isomorphic by a linear coordinate change and so are $\mathbb{Z}_4[x]/(h_j[x])$. Moreover, consider the homomorphism $\sigma : \mathbb{Z}_4[x]/(x^3+x+1) \to \mathbb{Z}_4[y]/(y^3-y+1)$ obtained by sending $\bar{x}$ to $\bar{y} + 2\bar{y}^2$. It is easy to see that σ is an isomorphism, and hence $R \simeq \mathbb{Z}_4[x]/(x^3 + x + 1)$. □

Corollary 3.3. *Let $(R, \mathfrak{m})$ be a finite local ring such that $|R| = 16$ and $|R/\mathfrak{m}| = 4$. If $char(R) = 4$, then $R \simeq \mathbb{Z}_4[x]/(x^2 + x + 1)$.*

Proof. We note that there is only one irreducible polynomial in $\mathbb{Z}_2[x]$ of degree 2, namely $x^2 + x + 1$. Therefore, applying Lemma 3.1 with $(q, p, d) = (4, 2, 2)$, we have that $R \simeq \mathbb{Z}_4[x]/(f(x))$, where $\overline{f(x)} = x^2+x+1$. Let $g_1(x) = x^2+x+1$, $g_2(x) = x^2+x+3$, $g_3(x) = x^2+3x+1$ and $g_4(x) = x^2+3x+3 \in \mathbb{Z}_4[x]$. Then $f(x)$ is one of these $g_i(x)$. Since $\mathbb{Z}_4[x]/(g_i(x))$ are isomorphic to each other, we conclude that $R \simeq \mathbb{Z}_4[x]/(x^2 + x + 1)$. □

We now recall a useful result from [20].

Theorem 3.4. *Let $(R, \mathfrak{m})$ be a finite local ring which is not a field. If $\gamma(\Gamma(R)) \leq 1$, then $|R/\mathfrak{m}| \leq 8$ and the following hold:*

(i) If $|R/\mathfrak{m}| = 8$, then $\mathfrak{m}^2 = 0$ and $|R| = 64$.

(ii) If $|R/\mathfrak{m}| = 7$, then $\mathfrak{m}^2 = 0$ and $|R| = 49$.

(iii) If $|R/\mathfrak{m}| = 5$, then $\mathfrak{m}^2 = 0$ and $|R| = 25$.

(iv) If $|R/\mathfrak{m}| = 4$, then $\mathfrak{m}^2 = 0$ and $|R| = 16$.

(v) If $|R/\mathfrak{m}| = 3$, then $\mathfrak{m}^3 = 0$ and $|R| \leq 27$.

(vi) If $|R/\mathfrak{m}| = 2$, then $\mathfrak{m}^5 = 0$ and $|R| \leq 32$.

Local rings with $\gamma(\Gamma(R)) \leq 1$

(3.5) From the above theorem we see that for a finite local ring $(R, \mathfrak{m})$ with $\gamma(\Gamma(R)) \leq 1$, the number of elements in its residue field is bounded above by 8. Hence, we shall analyze finite local rings by considering the cardinality of $R/\mathfrak{m}$.

$|R/\mathfrak{m}| = 8$.

From Theorem 3.4, we have that $\mathfrak{m}^2 = 0$ and $|R| = 64$. This implies that $|\mathfrak{m}| = 8$ and $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 1$ so that $\mathfrak{m} = (a)$ for every nonzero $a \in \mathfrak{m}$ and that $V(\Gamma(R)) = \mathfrak{m} - \{0\}$. Therefore $\Gamma(R) \simeq K_7$ and then $\gamma(\Gamma(R)) = 1$ by Lemma 1.1. We note that if $char(R) \geq 8$, then $\mathfrak{m} = (2)$. This implies that $char(R) = 4$, a contradiction. Thus, $char(R) \leq 4$.

(i) If $char(R) = 2$, then R is equi-characteristic, so that R contains the field $\mathbb{F}_8$. Since $\mathfrak{m}$ is principal, it then follows that $R = \mathbb{F}_8[\mathfrak{m}] \simeq \mathbb{F}_8[x]/(x^2)$.

(ii) If $char(R) = 4$, then from Corollary 3.2 we have that $R \simeq \mathbb{Z}_4[x]/(x^3 + x + 1)$.

$|R/\mathfrak{m}| = 7$.

From Theorem 3.4, we have that $\mathfrak{m}^2 = 0$ and $|R| = 49$. This implies that $|\mathfrak{m}| = 7$ and $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 1$, so that $\mathfrak{m} = (a)$ for every nonzero $a \in \mathfrak{m}$. We note that since $\mathfrak{m}^2 = 0$, $\Gamma(R) \simeq K_6$, so that $\gamma(\Gamma(R)) = \gamma(K_6) = 1$ by Lemma 1.1. Since $char(R)$ divides $|R|$, we have the following two cases to consider.

(i) If $char(R) = 7$, then $\mathbb{Z}_7 \subseteq R$. Since $\mathfrak{m}$ is principal, $R = \mathbb{Z}_7[\mathfrak{m}] \simeq \mathbb{Z}_7[x]/(x^2)$.

(ii) If $char(R) = 49$, then $\mathbb{Z}_{49} \subseteq R$. As $|R| = 49$, we conclude that $R = \mathbb{Z}_{49}$.

$|R/\mathfrak{m}| = 5$.

From Theorem 3.4, we have that $\mathfrak{m}^2 = 0$ and $|R| = 25$. This implies that $|\mathfrak{m}| = 5$ and $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 1$, so that $\mathfrak{m} = (a)$ for every nonzero $a \in \mathfrak{m}$. Since $\mathfrak{m}^2 = 0$, we have $\Gamma(R) \simeq K_4$ and thus $\gamma(\Gamma(R))$ is planar by Lemma 1.1.

(i) If $char(R) = 5$, then $R = \mathbb{Z}_5[\mathfrak{m}] \simeq \mathbb{Z}_5[x]/(x^2)$.

(ii) If $char(R) = 25$, then $\mathbb{Z}_{25} \subseteq R$ and thus $R = \mathbb{Z}_{25}$.

$|R/\mathfrak{m}| = 4$.

From Theorem 3.4, we have that $\mathfrak{m}^2 = 0$ and $|R| = 16$. This implies that $|\mathfrak{m}| = 4$ and $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 1$ so that $\mathfrak{m} = (a)$ for every nonzero $a \in \mathfrak{m}$. Since $\mathfrak{m}^2 = 0$, we have $\Gamma(R) \simeq K_3$, so that $\gamma(\Gamma(R))$ is planar. If $char(R) \geq 8$, then $\mathfrak{m} = (2)$, so that $char(R) = 4$, a contradiction. Thus $char(R) = 2$ or $char(R) = 4$.

(i) If $char(R) = 2$, then R is equi-characteristic, so that R contains the field $\mathbb{F}_4$. Since $\mathfrak{m}$ is principal, it follows that $R = \mathbb{F}_4[\mathfrak{m}] \simeq \mathbb{F}_4[x]/(x^2)$.

(ii) If $char(R) = 4$, then $R \simeq \mathbb{Z}_4[x]/(x^2 + x + 1)$ by Corollary 3.3.

$|R/\mathfrak{m}| = 3$.

From Theorem 3.4, we have that $\mathfrak{m}^3 = 0$ and $|R| \leq 27$, and therefore $|R| = 9$ or $|R| = 27$. We consider each case.

If $|R| = 9$, then $\mathfrak{m}^2 = 0$, so that $\Gamma(R) \simeq K_2$, which is planar. As in previous cases, we see that $R \simeq \mathbb{F}_3[x]/(x^2)$ or $R \simeq \mathbb{Z}_9$.

If $|R| = 27$, then $|\mathfrak{m}| = 9$. We note that $\mathfrak{m}^2 \neq 0$, for otherwise $\Gamma(R) \simeq K_8$ which implies $\gamma(\Gamma(R)) = 2$, a contradiction. It then follows from Proposition 2.4 and Example 2.5 that $\Gamma(R) \simeq \Gamma(\mathbb{Z}_{27}) \simeq G_1$, as shown in Figure 4-1.

(i) If $char(R) = 3$, then $R = \mathbb{F}_3[\mathfrak{m}] \simeq \mathbb{F}_3[x]/(x^3)$.

(ii) If $char(R) = 9$, then $3 \in \mathfrak{m}^2$. Otherwise $\mathfrak{m} = (3)$ and this implies that $\mathfrak{m}^2 = 0$, a contradiction. Suppose that $\mathfrak{m} = (a)$ for some $a \in \mathfrak{m} - \mathfrak{m}^2$, so that either $3 = a^2$ or $3 = -a^2$. Since $\mathfrak{m}$ is principal, $R = \mathbb{Z}_9[a]$ and R is isomorphic to a quotient ring of either $\mathbb{Z}_9[x]/(x^3, x^2 - 3)$ or $\mathbb{Z}_9[x]/(x^3, x^2 + 3)$. However, both rings above have 27 elements. Thus we conclude that $R \simeq \mathbb{Z}_9[x]/(x^3, x^2 - 3)$ or $R \simeq \mathbb{Z}_9[x]/(x^3, x^2 + 3)$.

(iii) If $char(R) = 27$, then $R = \mathbb{Z}_{27}$.

$|R/\mathfrak{m}| = 2$.

From Theorem 3.4, we have that $\mathfrak{m}^5 = 0$ and $|R| \leq 32$, and thus $|R| = 4, 8, 16$, or 32. As before, we consider cases accordingly.

If $|R| = 4$, then $\mathfrak{m}^2 = 0$, so that $\Gamma(R)$ consists of a single vertex. In this case, it is easy to see that $R \simeq \mathbb{Z}_4$ or $R \simeq \mathbb{Z}_2[x]/(x^2)$.

Assume that $|R| = 8$ and $\mathfrak{m}^2 = 0$. Then $|\mathfrak{m}| = 4$ and $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 2$. Therefore $\Gamma(R) \simeq K_3$, which is a planar graph.

(i) If $char(R) = 2$, then $R = \mathbb{Z}_2[\mathfrak{m}] \simeq \mathbb{Z}_2[x, y]/(x^2, xy, y^2)$.

(ii) If $char(R) = 4$, then $\mathfrak{m} = (2, a)$. We note that $2a = a^2 = 0$. Hence R is isomorphic to a quotient ring of $\mathbb{Z}_4[x]/(x^2, 2x)$. Since there are 8 elements in $\mathbb{Z}_4[x]/(x^2, 2x)$, we conclude that $R \simeq \mathbb{Z}_4[x]/(x^2, 2x)$.

Assume that $|R| = 8$ and $\mathfrak{m}^2 \neq 0$. By Proposition 2.4, we see that $\Gamma(R) \simeq \Gamma(\mathbb{Z}_8) \simeq P_3$, the path on 3 vertices.

(i) If $char(R) = 2$, then R contains the field $\mathbb{Z}_2$, so that $R \simeq \mathbb{Z}_2[x]/(x^3)$.

(ii) If $char(R) = 4$, then $\mathfrak{m}^2 = \{0, 2\}$. Otherwise $\mathfrak{m} = (2)$ and this implies $\mathfrak{m}^2 = 0$, which is a contradiction. Therefore $\mathfrak{m} = (a)$ with $a^2 - 2 = a^3 = 0$. It follows that R is isomorphic to a quotient ring of $\mathbb{Z}_4[x]/(x^3, x^2 - 2)$. However, $\mathbb{Z}_4[x]/(x^3, x^2 - 2)$ has 8 elements, so that $R \simeq \mathbb{Z}_4[x]/(x^3, x^2 - 2)$.

(iii) If $char(R) = 8$, then $R = \mathbb{Z}_8$.

Assume that $|R| = 16$ and $\mathfrak{m}^2 = 0$. It follows that $|\mathfrak{m}| = 8$ and $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 3$. Therefore, $\Gamma(R) \simeq K_7$ and thus $\gamma(\Gamma(R)) = \gamma(K_7) = 1$ by Lemma 1.1.

(i) If $char(R) = 2$, then $R = \mathbb{Z}_2[\mathfrak{m}] \simeq \mathbb{Z}_2[x, y, z]/(x, y, z)^2$.

(ii) If $char(R) = 4$, then $\mathfrak{m} = (2, a, b)$, so that $R \simeq \mathbb{Z}_4[x, y]/(x^2, xy, y^2, 2x, 2y)$.

Assume that $|R| = 16$, $\mathfrak{m}^2 \neq 0$, and $\mathfrak{m}^3 = 0$. If $|\mathfrak{m}^2| = 4$, then $\mathfrak{m}$ is principal and $|\mathfrak{m}^3| = 2$, which contradicts the assumption $\mathfrak{m}^3 = 0$. Therefore $|\mathfrak{m}^2| = 2$ and $\dim_{R/\mathfrak{m}} \mathfrak{m}/\mathfrak{m}^2 = 2$.

(i) Suppose $char(R) = 2$. Let $\mathfrak{m} = (a, b)$ for some $a, b \in \mathfrak{m}$. If $a^2 \neq 0$, then we may replace b if necessary and assume that $ab = 0$ as a^2 is a basis element of $\mathfrak{m}^2/\mathfrak{m}^3$. We note that $b^2 = 0$ or $b^2 = a^2$. Therefore R is isomorphic to either $\mathbb{Z}_2[x, y]/(x^3, xy, y^2)$ or $\mathbb{Z}_2[x, y]/(x^3, xy, y^2 - x^2)$. However, both rings in question have 16 elements, so that either $R \simeq \mathbb{Z}_2[x, y]/(x^3, xy, y^2)$ or $R \simeq \mathbb{Z}_2[x, y]/(x^3, xy, y^2 - x^2)$. In the first case, $\Gamma(R) \simeq K_{1,1,1,4}$ and thus $\gamma(\Gamma(R)) = 1$ by Example 2.9. In the other case, $\Gamma(R)$ is planar by Example 2.6. If $a^2 = b^2 = 0$, then R is isomorphic to $\mathbb{Z}_2[x, y]/(x^2, y^2)$. Again, since $\mathbb{Z}_2[x, y]/(x^2, y^2)$ has 16 elements, we conclude that $R \simeq \mathbb{Z}_2[x, y]/(x^2, y^2)$, and thus $\Gamma(R)$ is planar by Example 2.7.

(ii) Suppose $char(R) = 4$. First, we assume that $2 \notin \mathfrak{m}^2$. Then $\mathfrak{m} = (2, a)$ for some $a \in \mathfrak{m}$. If $a^2 \neq 0$, then $2a = 0$ or $2a = a^2$. By counting, we see that $R \simeq \mathbb{Z}_4[x]/(x^3, 2x)$ or $R \simeq \mathbb{Z}_4[x]/(x^3, x^2 - 2x)$. In the first case, $\gamma(\Gamma(R)) = 1$ by Example 2.9 and in the second case, $\Gamma(R)$ is planar by Example 2.6. If $a^2 = 0$, then $R \simeq \mathbb{Z}_4[x]/(x^2)$, which is planar by Example 2.7. Next, we assume that $2 \in \mathfrak{m}^2$. Let $\mathfrak{m} = (a, b)$ for some $a, b \in \mathfrak{m}$. From the remarks in (i), we see that if $a^2 \neq 0$, then we may assume that $ab = 0$ which implies that either $b^2 = 0$ or $b^2 = 2$. On the other hand, if $a^2 = b^2 = 0$, then $ab = 2$ as $\mathfrak{m}^2 \neq 0$. By counting, we conclude that $R \simeq \mathbb{Z}_4[x, y]/(x^3, x^2 - 2, xy, y^2)$ or $R \simeq \mathbb{Z}_4[x, y]/(x^3, x^2 - 2, xy, y^2 - 2, y^3)$ or $R \simeq \mathbb{Z}_4[x, y]/(x^2, xy - 2, y^2)$. In the first case, $\gamma(\Gamma(R)) = 1$ by Example 2.9. In the other two cases, we conclude $\Gamma(R)$ is planar by Examples 2.6 and 2.7 respectively.

(iii) Suppose $char(R) = 8$. We note that $\mathfrak{m} = (2, a)$ for some $a \in \mathfrak{m}$. For otherwise $2 \in \mathfrak{m}^2$, which implies $4 \in \mathfrak{m}^4 \neq 0$, a contradiction. Since 4 is the only nonzero element in $\mathfrak{m}^2$, we may replace a by $a - 2$ if necessary and assume that $2a = 0$. We note that $a^2 = 0$ or $a^2 - 4$, so that $R \simeq \mathbb{Z}_8[x]/(x^2, 2x)$ or $R \simeq \mathbb{Z}_8[x]/(x^2 - 4, 2x)$. In the first case, $\gamma(\Gamma(R)) = 1$ by Example 2.9 and $\Gamma(R)$ is planar in the second case by Example 2.6.

Assume that $|R| = 16$ and $\mathfrak{m}^3 \neq 0$. By Proposition 2.4, we see that $\mathfrak{m}$ is principal and $\Gamma(R) \simeq \Gamma(\mathbb{Z}_{16})$, which is planar by Example 2.5. As before, we now consider cases depending on the characteristic of R.

(i) Suppose $char(R) = 2$. In this case, $R \simeq \mathbb{Z}_2[x]/(x^4)$.

(ii) Suppose $char(R) = 4$. We note that $2 \in \mathfrak{m}^2$, as otherwise $\mathfrak{m} = (2)$, and then $\mathfrak{m}^2 = 0$, a contradiction. Let $\mathfrak{m} = (a)$ for some $a \in \mathfrak{m}$. Since $2a^2 = 0$, we see that $2 = a^2$ or $2 = a^3$ or $2 = a^2 + a^3$. This implies that $R \simeq \mathbb{Z}_4[x]/(x^2 - 2, x^4)$ or $R \simeq \mathbb{Z}_4[x]/(x^3 - 2, x^4)$ or $R \simeq \mathbb{Z}_4[x]/(x^3 + x^2 - 2, x^4)$.

(iii) Suppose $char(R) = 8$. We note that either $\mathfrak{m} = (2)$ or $2 \in \mathfrak{m}^2$, so that $\mathfrak{m}^3 = 0$ or $4 \in \mathfrak{m}^4 \neq 0$, both of which are impossible.

(iv) Suppose $char(R) = 16$. It is clear that $R = \mathbb{Z}_{16}$.

Assume that $|R| = 32$. We note that $\mathfrak{m}^2 \neq 0$. Otherwise $\Gamma(R) \simeq K_{15}$. This implies that $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.1, which is a contradiction. Hence $|\mathfrak{m}^2| = 2, 4$, or 8. However, if $|\mathfrak{m}^2| = 2$ or 4, then $\gamma(\Gamma(R)) \geq 2$ by Proposition 2.1 and Proposition 2.2. Therefore, $|\mathfrak{m}^2| = 8$, which implies that $\mathfrak{m}$ is principal and so is $\mathfrak{m}^k$ for each $k \in \mathbb{N}$. This implies that $\mathfrak{m}^4 \neq 0$. Thus, $\Gamma(R) \simeq \Gamma(\mathbb{Z}_{32})$ by Proposition 2.4 and $\gamma(\Gamma(R)) = 1$ by [20, Example 2.4].

(i) Suppose $char(R) = 2$. In this case, $R \simeq \mathbb{Z}_2[x]/(x^5)$.

(ii) Suppose $char(R) = 4$. We note that $2 \in \mathfrak{m}^3$, as otherwise either $\mathfrak{m} = (2)$ or $\mathfrak{m}^2 = (2)$ and both imply that $\mathfrak{m}^4 = 0$, a contradiction. Suppose that $\mathfrak{m} = (a)$. Since $2a^2 = 0$, we see that $2 = a^3$ or $2 = a^3 + a^4 = (a + a^2)^3$ or $2 = a^4$. It follows that $R \simeq \mathbb{Z}_4[x]/(x^3 - 2, x^5) \simeq \mathbb{Z}_4[x]/((x + x^2)^3 - 2, x^5)$ or $R \simeq \mathbb{Z}_4[x]/(x^4 - 2, x^5)$.

(iii) Suppose $char(R) = 8$. If $\mathfrak{m} = (2)$ or $2 \in \mathfrak{m}^3$, then $\mathfrak{m}^3 = 0$ or $4 \in \mathfrak{m}^6 \neq 0$, which are impossible. Thus, $\mathfrak{m}^2 = (2)$. Let $\mathfrak{m} = (a)$ for some $a \in \mathfrak{m}$. Since $2a^2 = 4 \in \mathfrak{m}^4$ and $4a = 0 \in \mathfrak{m}^5$, we have that either $2 = a^2$, $2 = -a^2 = (a + 2)^2$, $2 = a^2 + 2a$, or $2 = -a^2 + 2a$. Accordingly, it follows that $R \simeq \mathbb{Z}_8[x]/(x^2 - 2, x^5) \simeq \mathbb{Z}_8[x]/((x + 2)^2 - 2, x^5)$ or $R \simeq \mathbb{Z}_8[x]/(x^2 + 2x - 2, x^5)$ or $R \simeq \mathbb{Z}_8[x]/(x^2 - 2x + 2, x^5)$.

(iv) Suppose $char(R) = 16$. We note either $\mathfrak{m} = (2)$ or $2 \in \mathfrak{m}^2$. This implies either $\mathfrak{m}^4 = 0$ or $8 \in \mathfrak{m}^6 \neq 0$, both of which are impossible.

(v) If $char(R) = 32$, it is clear that $R = \mathbb{Z}_{32}$.

Summarizing the above, we obtain the following theorems which completely classify those local rings with a toroidal zero-divisor graph. We are primarily concerned with the genus one case, but our analysis recovers the following result from [18]. Note that in that paper, some of the ideals are chosen with different generators. For instance, in that paper the rings $\mathbb{Z}_9[x]/(x^2 - 3, 3x)$ and $\mathbb{Z}_9[x]/(x^2 - 6, 3x)$ are listed as having planar zero-divisor graphs. In our paper, we use different generators for the ideals in question.

Theorem 3.5.1. *Let $(R, \mathfrak{m})$ be a finite local ring which is not a field. Then $\Gamma(R)$ is planar if and only if R is isomorphic to one of the following 29 rings.*

$\mathbb{Z}_4$, $\mathbb{Z}_8$, $\mathbb{Z}_9$, $\mathbb{Z}_{16}$, $\mathbb{Z}_{25}$, $\mathbb{Z}_{27}$, $\frac{\mathbb{Z}_2[x]}{(x^2)}$, $\frac{\mathbb{Z}_2[x]}{(x^3)}$, $\frac{\mathbb{Z}_2[x]}{(x^4)}$, $\frac{\mathbb{Z}_2[x,y]}{(x^2,y^2)}$ $\frac{\mathbb{Z}_2[x,y]}{(x^2,xy,y^2)}$, $\frac{\mathbb{Z}_2[x,y]}{(x^3,xy,y^2-x^2)}$, $\frac{\mathbb{F}_4[x]}{(x^2)}$, $\frac{\mathbb{Z}_3[x]}{(x^2)}$, $\frac{\mathbb{Z}_3[x]}{(x^3)}$, $\frac{\mathbb{Z}_4[x]}{(x^2)}$, $\frac{\mathbb{Z}_4[x]}{(x^2+x+1)}$, $\frac{\mathbb{Z}_4[x]}{(2x,x^2)}$, $\frac{\mathbb{Z}_4[x]}{(x^2-2,x^4)}$, $\frac{\mathbb{Z}_4[x]}{(x^3-2,x^4)}$, $\frac{\mathbb{Z}_4[x]}{(x^2-2,x^3)}$, $\frac{\mathbb{Z}_4[x]}{(x^3,x^2-2x)}$, $\frac{\mathbb{Z}_4[x]}{(x^3+x^2-2,x^4)}$, $\frac{\mathbb{Z}_4[x,y]}{(x^2,y^2,xy-2)}$, $\frac{\mathbb{Z}_4[x,y]}{(x^3,x^2-2,xy,y^2-2,y^3)}$, $\frac{\mathbb{Z}_5[x]}{(x^2)}$, $\frac{\mathbb{Z}_8[x]}{(x^2-4,2x)}$, $\frac{\mathbb{Z}_9[x]}{(x^2-3,x^3)}$, $\frac{\mathbb{Z}_9[x]}{(x^2+3,x^3)}$.

Theorem 3.5.2. *Let $(R, \mathfrak{m})$ be a finite local ring which is not a field. Then $\gamma(\Gamma(R)) = 1$ if and only if R is isomorphic to one of the following 17 rings.*

$\mathbb{Z}_{32}$, $\mathbb{Z}_{49}$, $\frac{\mathbb{Z}_2[x]}{(x^5)}$, $\frac{\mathbb{F}_8[x]}{(x^2)}$, $\frac{\mathbb{Z}_2[x,y]}{(x^3,xy,y^2)}$, $\frac{\mathbb{Z}_2[x,y,z]}{(x,y,z)^2}$, $\frac{\mathbb{Z}_4[x]}{(x^3+x+1)}$, $\frac{\mathbb{Z}_4[x]}{(x^3-2,x^5)}$, $\frac{\mathbb{Z}_4[x]}{(x^4-2,x^5)}$, $\frac{\mathbb{Z}_4[x,y]}{(x^3,x^2-2,xy,y^2)}$, $\frac{\mathbb{Z}_4[x]}{(x^3,2x)}$, $\frac{\mathbb{Z}_4[x,y]}{(2x,2y,x^2,xy,y^2)}$, $\frac{\mathbb{Z}_7[x]}{(x^2)}$, $\frac{\mathbb{Z}_8[x]}{(x^2,2x)}$, $\frac{\mathbb{Z}_8[x]}{(x^2-2,x^5)}$, $\frac{\mathbb{Z}_8[x]}{(x^2+2x-2,x^5)}$, $\frac{\mathbb{Z}_8[x]}{(x^2-2x+2,x^5)}$.

Non-local rings with $\gamma(\Gamma(R)) \leq 1$

(3.6) Since a finite ring is Artinian, it is isomorphic to a finite direct product of Artinian local rings (see [1, Theorem 8.7]). Thus, the number of maximal ideals of R is simply the number of components of R. From [20, Lemma 3.1], we know that for a finite ring R, $|Spec(R)| \geq 5$ implies that $\gamma(\Gamma(R)) \geq 2$. Thus, to find all rings R with $\gamma(\Gamma(R)) \leq 1$, we need only consider all R with $|Spec(R)| \leq 4$. We thus consider cases according to the cardinality of $Spec(R)$. As before, we will recover known results about the planar case as part of our analysis.

Let $|Spec(R)| = 2$.

By assumption, $R \simeq R_1 \times R_2$ where both $(R_1, \mathfrak{m}_1)$ and $(R_2, \mathfrak{m}_2)$ are finite local rings with $\gamma(\Gamma(R_i) \leq 1$. If R_1 and R_2 are both fields, then $\Gamma(R) \simeq K_{m,n}$, where $m = |R_1| - 1$ and $n = |R_2| - 1$. Therefore, by Lemma 1.2, $\Gamma(R)$ is planar if and only if $R \simeq \mathbb{Z}_2 \times \mathbb{F}_q$ or $R \simeq \mathbb{Z}_3 \times \mathbb{F}_q$, where $\mathbb{F}_q$ is a finite field with q elements. Further, $\gamma(\Gamma(R)) = 1$ if and only if $R \simeq \mathbb{F}_4 \times \mathbb{F}_4$, $R \simeq \mathbb{F}_4 \times \mathbb{Z}_5$, $R \simeq \mathbb{F}_4 \times \mathbb{Z}_7$, or $R \simeq \mathbb{Z}_5 \times \mathbb{Z}_5$. Thus, we may assume that at least one of the R_i is not a field. We proceed by considering the pair $(|R_1|, |R_2|)$. Without loss, we assume that $|R_1| \leq |R_2|$.

Case 1 : $|R_1| = 2$. In this case, $R_1 \simeq \mathbb{Z}_2$. Since R_2 is not a field, we have $|R_2/\mathfrak{m}_2| \leq 8$ by Theorem 3.4, and also we see that if $|R_2/\mathfrak{m}_2| \geq 5$, then $\mathfrak{m}_2^2 = 0$. Let $a_1, \ldots, a_4$ be distinct nonzero elements in $\mathfrak{m}_2$ and let $u_i = (0, a_i)$, $v_i = (1, a_i)$ for $i = 1, \ldots, 4$, and $v_5 = (1, 0)$. Then $u_i \cdot v_j = 0$ for every i, j, so that $K_{4,5} \subseteq \Gamma(\mathbb{Z}_2 \times R_2)$. It follows that $\gamma(\Gamma(\mathbb{Z}_2 \times R_2)) \geq 2$. We may thus assume that $|R_2/\mathfrak{m}_2| \leq 4$. We now proceed by cases, according to the cardinality of $|R_2/\mathfrak{m}_2|$.

Assume that $|R_2/\mathfrak{m}_2| = 4$. Then we have that $|\mathfrak{m}_2| = 4$, $\mathfrak{m}_2^2 = 0$, and that $\mathfrak{m}_2$ is principal by Theorem 3.4. Therefore, $\gamma(\Gamma(\mathbb{Z}_2 \times R_2)) = 1$ by Proposition 2.11. We see from the discussion in (3.5) that $R_2 \simeq \dfrac{\mathbb{F}_4[x]}{(x^2)}$ or $\dfrac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}$, so that $R \simeq \mathbb{Z}_2 \times \dfrac{\mathbb{F}_4[x]}{(x^2)}$ or $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}$ in this case.

Assume that $|R_2/\mathfrak{m}_2| = 3$. Then $\mathfrak{m}_2^3 = 0$ and $|R_2| \leq 27$ by Theorem 3.4.

(i) If $|R_2| = 3$, then $R_2 \simeq \mathbb{Z}_3$ and $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3)$ is planar by [4, Theorem 5.1].

(ii) If $|R_2| = 9$, then $R_2 \simeq \mathbb{Z}_9$ or $R_2 \simeq \dfrac{\mathbb{Z}_3[x]}{(x^2)}$. However, $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_9) \simeq \Gamma(\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_3[x]}{(x^2)})$ by Proposition 2.4 and $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_9)$ is planar by [4, Theorem 5.1], so $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_9$ or $R \simeq \mathbb{Z}_2 \times \dfrac{\mathbb{Z}_3[x]}{(x^2)}$ in this case.

(iii) If $|R_2| = 27$, then $\mathfrak{m}_2^2 \neq 0$ from (3.5), so that $\Gamma(\mathbb{Z}_2 \times R_2) \simeq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_{27})$ by Proposition 2.4. However, $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_{27})) \geq 2$ by [20, Theorem 4.5].

Now, assume that $|R_2/\mathfrak{m}_2| = 2$. Then $\mathfrak{m}_2^5 = 0$ and $|R_2| \leq 32$ by Theorem 3.4.

(i) If $|R_2| = 4$, then $R_2 \simeq \mathbb{Z}_4$ or $R_2 \simeq \dfrac{\mathbb{Z}_2[x]}{(x^2)}$. However, since $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_4) \simeq \Gamma(\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)})$ and $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_4)$ is planar by [4, Theorem 5.1], we see that $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_4$ or $R \simeq \mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$.

(ii) Suppose that $|R_2| = 8$ and $\mathfrak{m}_2^2 = 0$. In this case, $|\mathfrak{m}_2| = 4$, so that $\gamma(\Gamma(\mathbb{Z}_2 \times R_2)) = 1$ by Proposition 2.11. Therefore, $R \simeq \mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x, y]}{(x^2, xy, y^2)}$, or $R \simeq \mathbb{Z}_2 \times \dfrac{\mathbb{Z}_4[x]}{(2x, x^2)}$.

(iii) Suppose that $|R_2| = 8$ and $\mathfrak{m}_2^2 \neq 0$. By Proposition 2.4 we have that $\Gamma(R_2) \simeq \Gamma(\mathbb{Z}_8)$ and that $\Gamma(\mathbb{Z}_2 \times R_2) \simeq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_8)$, which is planar by [4, Theorem 5.1]. Thus, $R \simeq \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^3)}$ or $R \simeq \mathbb{Z}_2 \times \frac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$ or $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_8$ by (3.5).

(iv) Suppose that $|R_2| = 16$. In this case, $\mathfrak{m}_2^3 \neq 0$, as otherwise $\gamma(\Gamma(R)) \geq 2$ by Proposition 2.3. Therefore, $\Gamma(R_2) \simeq \Gamma(\mathbb{Z}_{16})$ and $\Gamma(R) \simeq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_{16})$ by Proposition 2.4. However, $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_{16})) \geq 2$ by [20, Example 2.5].

(v) Finally, suppose that $|R_2| = 32$. Then $\mathfrak{m}_2^4 \neq 0$ and $\mathfrak{m}_2^5 = 0$ from the discussion in (3.5). Thus, $\Gamma(R_2) \simeq \Gamma(\mathbb{Z}_{32})$ and $\Gamma(R) \simeq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_{32})$. However, $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_{32})) \geq 2$ by [20, Example 2.5].

Case 2 : $|R_1| = 3$. In this case, $R_1 \simeq \mathbb{Z}_3$. Since R_2 is not a field, $|R_2/\mathfrak{m}_2| \leq 8$ by Theorem 3.4. If $|R_2/\mathfrak{m}_2| \geq 4$, then $\mathfrak{m}_2^2 = 0$ by Theorem 3.4. Let a_1, a_2, a_3 be distinct nonzero elements in $\mathfrak{m}_2$ and let $u_i = (0, a_i)$, $v_i = (1, a_i)$, $v_{i+3} = (2, a_i)$ for $i = 1, 2, 3$, and let $v_7 = (1, 0)$. Then $u_i \cdot v_j = 0$ for every i, j so that $K_{3,7} \subseteq \Gamma(\mathbb{Z}_3 \times R_2)$ and therefore $\gamma(\Gamma(\mathbb{Z}_3 \times R_2)) \geq 2$. We may thus assume henceforth that $|R_2/\mathfrak{m}_2| \leq 3$.

Assume that $|R_2/\mathfrak{m}_2| = 3$. Then $\mathfrak{m}_2^3 = 0$ and $|R_2| \leq 27$ by Theorem 3.4.

(i) If $|R_2| = 9$, then $R_2 \simeq \mathbb{Z}_9$ or $\frac{\mathbb{Z}_3[x]}{(x^2)}$. We note that $\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_9) \simeq \Gamma(\mathbb{Z}_3 \times \frac{\mathbb{Z}_3[x]}{(x^2)})$ by Proposition 2.4 and $\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_9)$ is planar by [4, Theorem 5.1]. Therefore, $R \simeq \mathbb{Z}_3 \times \mathbb{Z}_9$ or $R \simeq \mathbb{Z}_3 \times \frac{\mathbb{Z}_3[x]}{(x^2)}$ in this case.

(ii) If $|R_2| = 27$, then $\mathfrak{m}_2^2 \neq 0$ by (3.5), so that $\Gamma(\mathbb{Z}_3 \times R_2) \simeq \Gamma(\mathbb{Z}_3 \times \mathbb{Z}_{27})$ by Proposition 2.4. However, in this case $\gamma(\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_{27})) \geq 2$.

Assume that $|R_2/\mathfrak{m}_2| = 2$. Then $\mathfrak{m}_2^5 = 0$ and $|R_2| \leq 32$ by Theorem 3.4.

(i) If $|R_2| = 4$, then $R_2 \simeq \mathbb{Z}_4$ or $\frac{\mathbb{Z}_2[x]}{(x^2)}$. We note that $\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_4) \simeq \Gamma(\mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^2)})$ and $\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_4)$ is planar by [4, Theorem 5.1], so $R \simeq \mathbb{Z}_3 \times \mathbb{Z}_4$ or $R \simeq \mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$ in this case.

(ii) Suppose that $|R_2| = 8$ and $\mathfrak{m}_2^2 = 0$. Let $G = \Gamma(\mathbb{Z}_3 \times R_2)$ and let a_1, a_2, a_3 be distinct nonzero elements in $\mathfrak{m}_2$. Let $u_i = (0, a_i)$, $v_i = (1, a_i)$ and $v_{i+3} = (2, a_i)$ for $i = 1, 2, 3$, and let $v_7 = (1, 0)$. We then see that $u_i \cdot v_j = 0$ for every i, j as $\mathfrak{m}_2^2 = 0$ by Theorem 3.4, so that $K_{3,7} \subseteq \Gamma(\mathbb{Z}_3 \times R_2)$. It then follows that $\gamma(\Gamma(\mathbb{Z}_3 \times R_2)) \geq 2$ by Lemma 1.2.

(iii) Suppose that $|R_2| = 8$ and $\mathfrak{m}_2^2 \neq 0$. Then $\Gamma(R_2) \simeq \Gamma(\mathbb{Z}_8)$ and $\Gamma(\mathbb{Z}_3 \times R_2) \simeq \Gamma(\mathbb{Z}_3 \times \mathbb{Z}_8)$ by Proposition 2.4. We note that $\gamma(\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_8)) = 1$ by [20, Theorem 4.5]. Thus either $R \simeq \mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^3)}$, $R \simeq \mathbb{Z}_3 \times \frac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$, or $R \simeq \mathbb{Z}_3 \times \mathbb{Z}_8$ from the discussion in (3.5).

(iv) If $|R_2| \geq 16$, then $\Gamma(\mathbb{Z}_2 \times R_2)$ is a subgraph of $\Gamma(R)$. However, we see from Case 1.C.(iv) that $\gamma(\Gamma(\mathbb{Z}_2 \times R_2)) \geq 2$.

Case 3 : $|R_1| = 4$. In this case, $R_1 \simeq \mathbb{F}_4$ or $\mathbb{Z}_4$ or $\frac{\mathbb{Z}_2[x]}{(x^2)}$. We note that if $|R_2| \geq 8$, then $K_{3,7} \subseteq \Gamma(R)$, and so $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.2. Therefore, $|R_2| = 4, 5$, or 7. Equivalently, $R_2 \simeq \mathbb{F}_4, \mathbb{Z}_4, \frac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_5$,

or $\mathbb{Z}_7$. If $R_1 \simeq \mathbb{F}_4$, then $R_2 \simeq \mathbb{Z}_4$ or $\frac{\mathbb{Z}_2[x]}{(x^2)}$ as R_2 is not a field. It is easy to see that the reduction of $\Gamma(\mathbb{F}_4 \times \mathbb{Z}_4)$ is isomorphic to $K_{3,3}$. Since $\gamma(K_{3,3}) = 1$ and $\Gamma(\mathbb{F}_4 \times \mathbb{Z}_4) \simeq \Gamma(\mathbb{F}_4 \times \frac{\mathbb{Z}_2[x]}{(x^2)})$, then $R \simeq \mathbb{F}_4 \times \mathbb{Z}_4$ or $\mathbb{F}_4 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$. On the other hand, if $R_1 \simeq \mathbb{Z}_4$ and $R_2 \simeq \mathbb{Z}_n$, then by [20, Theorem 3.5] we see that $\Gamma(\mathbb{Z}_4 \times \mathbb{Z}_n)$ is of genus one for $n = 4, 5$, and 7. By Proposition 2.4, we have $\Gamma(\mathbb{Z}_4 \times S) \simeq \Gamma(\frac{\mathbb{Z}_2[x]}{(x^2)} \times S)$ for any ring S. We thus conclude that $R \simeq \mathbb{Z}_4 \times \mathbb{Z}_4$, $\mathbb{Z}_4 \times \mathbb{Z}_5$, $\mathbb{Z}_4 \times \mathbb{Z}_7$, $\mathbb{Z}_4 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_5 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_7 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$, or $\frac{\mathbb{Z}_2[x]}{(x^2)} \times \frac{\mathbb{Z}_2[x]}{(x^2)}$.

Case 4 : $|R_1| \geq 5$. Then $|R_2| \geq 8$ as $|R_2| \geq |R_1|$ by assumption. Therefore, $K_{4,7} \subseteq \Gamma(R)$ and then $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.2.

Let $|Spec(R)| = 3$.

Since $|Spec(R)| = 3$, there are three finite local rings $(R_i, \mathfrak{m}_i)$, $i = 1, 2, 3$ such that $R \simeq R_1 \times R_2 \times R_3$. We proceed by considering the triple $(|R_1|, |R_2|, |R_3|)$. Without loss we may assume that $|R_1| \leq |R_2| \leq |R_3|$, and we consider the possibilities for such a triple.

Suppose $|R_1| \geq 3$. In this case, $R_i \geq 3$ for each i. If $|R_3| \geq 4$, then $K_{3,8} \subseteq \Gamma(R)$, which implies that $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.2, a contradiction. Therefore, we may assume that $|R_3| \leq 3$. Then $|R_i| = 3$ for each i. Note that in this case, $\gamma(\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3)) = 1$ by [20, Example 2.4].

Suppose $|R_1| = 2$ and $|R_2| \geq 3$. In this case, if $|R_2| \geq 4$ or $|R_3| \geq 5$, then $K_{3,7}$ or $K_{4,5}$ is contained in $\Gamma(R)$, so that $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.2. Therefore, we may assume that $(|R_2|, |R_3|) = (3, 3)$ or $(3, 4)$. If $(|R_2|, |R_3|) = (3, 3)$, then $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3$. In this case we have $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3)) = 1$ by [4, Theorem 5.1] and $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3)$ is a subgraph of $\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3)$. Now, if $(|R_2|, |R_3|) = (3, 4)$ then $R_2 = \mathbb{Z}_3$ and $R_3 = \mathbb{F}_4$, $\mathbb{Z}_4$, or $\frac{\mathbb{Z}_2[x]}{(x^2)}$. Example 2.10 shows that $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{F}_4)) = 1$. However, by [20, Example 2.5], we have $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4)) \geq 2$ and $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^2)})) \geq 2$ as well, since $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^2)}) \simeq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_4)$ by Proposition 2.4. Therefore we must have $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3$ or $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{F}_4$.

Suppose $|R_1| = 2$ and $|R_2| = 2$. In this case, if $|R_3| \geq 8$, then $K_{3,7}$ is contained in $\Gamma(R)$ which forces $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.2. Therefore, we may assume that $|R_3| \leq 7$. Note that $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_n$ is planar when $n = 2$ or 3 by [4, Theorem 5.1] and $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_n)) = 1$ when $n = 5$ or 7 by [20, Theorem 3.5]. Thus, it remains only to examine the case where $|R_3| = 4$, that is where $R_3 = \mathbb{F}_4$, $\mathbb{Z}_4$, or $\frac{\mathbb{Z}_2[x]}{(x^2)}$. We note that $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}) \simeq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4)$ by Proposition 2.4 and $K_{3,3} \subseteq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{F}_4) \subseteq \Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4)$ by Remark 2.12. Since $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4)) = 1$ by [20, Example 2.4], we see that $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4)) = \gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}) = \gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{F}_4)) = 1$. Therefore, $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{F}_4$ or

$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$ or $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$ in this case.

Suppose $|Spec(R)| = 4$**.**

We assume that $R \simeq R_1 \times R_2 \times R_3 \times R_4$ where each $(R_i, \mathfrak{m}_i)$ is a finite local ring for $i = 1, 2, 3, 4$. As before we also assume that $|R_1| \leq |R_2| \leq |R_3| \leq |R_4|$.

Case 1 : $|R_3| \geq 3$. In this case, certainly $|R_4| \geq 3$. Therefore, $K_{3,8} \subseteq \Gamma(R)$, and then $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.2.

Case 2 : $|R_1| = |R_2| = |R_3| = 2$. In this case, if $|R_4| \geq 4$, then $K_{3,7} \subseteq \Gamma(R)$, so that $\gamma(\Gamma(R)) \geq 2$ by Lemma 1.2, a contradiction. Therefore, $|R_4| \leq 3$, so that $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ or $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3$. However, $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2)) = 1$ and $\gamma(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) \geq 2$ by [20, Example 2.5], so that $R \simeq \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ in this case.

We summarize the above work to obtain two theorems. These two results provide a complete list of non-local rings whose zero-divisor graphs have genera at most one. The planar case has been discussed previously in [2], [7] and in [18]. Also, the next theorem appears in [18]; we state the result since it follows immediately from our analysis. As before, note that in [18] different generators are chosen for some of the ideals.

Theorem 3.6.1. *Let R be a finite ring which is not local; then $\Gamma(R)$ is planar if and only if R is isomorphic to one of the following 15 types of rings.*
$\mathbb{Z}_2 \times \mathbb{F}_q$, $\mathbb{Z}_3 \times \mathbb{F}_q$, $\mathbb{Z}_2 \times \mathbb{Z}_9$, $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_3[x]}{(x^2)}$, $\mathbb{Z}_2 \times \mathbb{Z}_4$, $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x]}{(x^3)}$, $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$, $\mathbb{Z}_2 \times \mathbb{Z}_8$, $\mathbb{Z}_3 \times \mathbb{Z}_9$, $\mathbb{Z}_3 \times \dfrac{\mathbb{Z}_3[x]}{(x^2)}$, $\mathbb{Z}_3 \times \mathbb{Z}_4$, $\mathbb{Z}_3 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3$.

Theorem 3.6.2. *Let R be a finite ring which is not local; then $\gamma(\Gamma(R)) = 1$ if and only if R is isomorphic to one of the following 29 rings.*
$\mathbb{F}_4 \times \mathbb{F}_4$, $\mathbb{F}_4 \times \mathbb{Z}_5$, $\mathbb{F}_4 \times \mathbb{Z}_7$, $\mathbb{Z}_5 \times \mathbb{Z}_5$, $\mathbb{Z}_2 \times \dfrac{\mathbb{F}_4[x]}{(x^2)}$, $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_4[x]}{(x^2+x+1)}$, $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x,y]}{(x^2, xy, y^2)}$, $\mathbb{Z}_2 \times \dfrac{\mathbb{Z}_4[x]}{(2x, x^2)}$, $\mathbb{Z}_3 \times \dfrac{\mathbb{Z}_2[x]}{(x^3)}$, $\mathbb{Z}_3 \times \dfrac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$, $\mathbb{Z}_3 \times \mathbb{Z}_8$, $\mathbb{Z}_4 \times \mathbb{F}_4$, $\mathbb{F}_4 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_4 \times \mathbb{Z}_4$, $\mathbb{Z}_4 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, $\dfrac{\mathbb{Z}_2[x]}{(x^2)} \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_4 \times \mathbb{Z}_5$, $\mathbb{Z}_5 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_4 \times \mathbb{Z}_7$, $\mathbb{Z}_7 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3$, $\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3$, $\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{F}_4$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_7$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{F}_4$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$, $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \dfrac{\mathbb{Z}_2[x]}{(x^2)}$, *and* $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$.

At the end of this paper, we summarize all the results obtained through the discussion in (3.5), (3.6) in four tables showed in the following pages.

Acknowledgement

The authors wish to express their deepest gratitude to the referee for careful reading of the paper, helpful comments and many valuable suggestions.

G_1, G_2, G_3, G_4 are graphs as shown in Figure 4-1, 4-2, 5-1, 5-2.

R	$\mathbb{Z}_4$	$\frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{Z}_9$	$\frac{\mathbb{Z}_3[x]}{(x^2)}$	$\mathbb{Z}_8$	$\frac{\mathbb{Z}_2[x]}{(x^3)}$	$\frac{\mathbb{Z}_4[x]}{(x^2-2,x^3)}$	$\frac{\mathbb{Z}_2[x,y]}{(x^2,xy,y^2)}$	$\frac{\mathbb{Z}_4[x]}{(2x,x^2)}$	$\frac{\mathbb{F}_4[x]}{(x^2)}$
$\|R\|$	4	4	9	9	8	8	8	8	8	16
$\|R/\mathfrak{m}\|$	2	2	3	3	2	2	2	2	2	4
$char(R)$	4	2	9	3	8	2	4	2	4	2
$\Gamma(R)$	Point	Point	K_2	K_2	P_3	P_3	P_3	K_3	K_3	K_3
$\|V(\Gamma(R))\|$	1	1	2	2	3	3	3	3	3	3
$\gamma(\Gamma(R))$	0	0	0	0	0	0	0	0	0	0

R	$\frac{\mathbb{Z}_4[x]}{(x^2+x+1)}$	$\mathbb{Z}_{25}$	$\frac{\mathbb{Z}_5[x]}{(x^2)}$	$\mathbb{Z}_{16}$	$\frac{\mathbb{Z}_2[x]}{(x^4)}$	$\frac{\mathbb{Z}_4[x]}{(x^2-2,x^4)}$	$\frac{\mathbb{Z}_4[x]}{(x^3-2,x^4)}$	$\frac{\mathbb{Z}_4[x]}{(x^3+x^2-2,x^4)}$	$\frac{\mathbb{Z}_2[x,y]}{(x^3,xy,y^2-x^2)}$	$\frac{\mathbb{Z}_4[x]}{(x^3,x^2-2x)}$
$\|R\|$	16	25	25	16	16	16	16	16	16	16
$\|R/\mathfrak{m}\|$	4	5	5	2	2	2	2	2	2	2
$char(R)$	4	25	5	16	2	4	4	4	2	4
$\Gamma(R)$	K_3	K_4	K_4	G_2	G_2	G_2	G_2	G_2	G_3	G_3
$\|V(\Gamma(R))\|$	3	4	4	7	7	7	7	7	7	7
$\gamma(\Gamma(R))$	0	0	0	0	0	0	0	0	0	0

R	$\frac{\mathbb{Z}_8[x]}{(x^2-4,2x)}$	$\frac{\mathbb{Z}_4[x,y]}{(x^3,x^2-2,xy,y^2-2,y^3)}$	$\frac{\mathbb{Z}_4[x]}{(x^2)}$	$\frac{\mathbb{Z}_4[x,y]}{(x^2,y^2,xy-2)}$	$\frac{\mathbb{Z}_2[x,y]}{(x^2,y^2)}$	$\mathbb{Z}_{27}$	$\frac{\mathbb{Z}_3[x]}{(x^3)}$	$\frac{\mathbb{Z}_9[x]}{(x^2-3,x^3)}$	$\frac{\mathbb{Z}_9[x]}{(x^2+3,x^3)}$
$\|R\|$	16	16	16	16	16	27	27	27	27
$\|R/\mathfrak{m}\|$	2	2	2	2	2	3	3	3	3
$char(R)$	8	4	4	4	2	27	3	9	9
$\Gamma(R)$	G_3	G_3	G_4	G_4	G_4	G_1	G_1	G_1	G_1
$\|V(\Gamma(R))\|$	7	7	7	7	7	8	8	8	8
$\gamma(\Gamma(R))$	0	0	0	0	0	0	0	0	0

Table 1. Local rings with planar zero-divisor graphs.

G_5 is the graph mentioned in Remark 2.8.

R	$\mathbb{Z}_{49}$	$\frac{\mathbb{Z}_7[x]}{(x^2)}$	$\frac{\mathbb{Z}_2[x,y]}{(x^3,xy,y^2)}$	$\frac{\mathbb{Z}_4[x]}{(x^3,2x)}$	$\frac{\mathbb{Z}_4[x,y]}{(x^3,x^2-2,xy,y^2)}$	$\frac{\mathbb{Z}_8[x]}{(x^2,2x)}$	$\frac{\mathbb{F}_8[x]}{(x^2)}$	$\frac{\mathbb{Z}_4[x]}{(x^3+x+1)}$	$\frac{\mathbb{Z}_4[x,y]}{(2x,2y,x^2,xy,y^2)}$	$\frac{\mathbb{Z}_2[x,y,z]}{(x,y,z)^2}$
$\lvert R\rvert$	49	49	16	16	16	16	64	64	16	16
$\lvert R/\mathfrak{m}\rvert$	7	7	2	2	2	2	8	8	2	2
$char(R)$	49	7	2	4	4	8	2	4	4	2
$\Gamma(R)$	K_6	K_6	$K_{1,1,1,4}$	$K_{1,1,1,4}$	$K_{1,1,1,4}$	$K_{1,1,1,4}$	K_7	K_7	K_7	K_7
$\lvert V(\Gamma(R))\rvert$	6	6	7	7	7	7	7	7	7	7
$\gamma(\Gamma(R))$	1	1	1	1	1	1	1	1	1	1

R	$\mathbb{Z}_{32}$	$\frac{\mathbb{Z}_2[x]}{(x^5)}$	$\frac{\mathbb{Z}_4[x]}{(x^3-2,x^5)}$	$\frac{\mathbb{Z}_4[x]}{(x^4-2,x^5)}$	$\frac{\mathbb{Z}_8[x]}{(x^2-2,x^5)}$	$\frac{\mathbb{Z}_8[x]}{(x^2-2x+2,x^5)}$	$\frac{\mathbb{Z}_8[x]}{(x^2+2x-2,x^5)}$
$\lvert R\rvert$	32	32	32	32	32	32	32
$\lvert R/\mathfrak{m}\rvert$	2	2	2	2	2	2	2
$char(R)$	32	2	4	4	8	8	8
$\Gamma(R)$	G_5	G_5	G_5	G_5	G_5	G_5	G_5
$\lvert V(\Gamma(R))\rvert$	15	15	15	15	15	15	15
$\gamma(\Gamma(R))$	1	1	1	1	1	1	1

Table 2. Local rings with toroidal zero-divisor graphs.

p is prime and $p \neq 2, 3$.

R	$\mathbb{Z}_2 \times \mathbb{F}_{p^n}$	$\mathbb{Z}_3 \times \mathbb{F}_{p^n}$	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$\mathbb{Z}_2 \times \mathbb{Z}_3$	$\mathbb{Z}_2 \times \mathbb{F}_4$	$\mathbb{Z}_3 \times \mathbb{Z}_3$	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$\mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{Z}_3 \times \mathbb{F}_4$	$\mathbb{Z}_3 \times \mathbb{Z}_4$
$\|R\|$	$2p^n$	$3p^n$	4	6	8	9	8	8	12	12
$\|Spec(R)\|$	2	2	2	2	2	2	2	2	2	2
$char(R)$	$2p$	$3p$	2	6	2	3	4	2	6	12
$\|V(\Gamma(R))\|$	p^n	$p^n + 1$	2	3	4	4	5	5	5	7
$\gamma(\Gamma(R))$	0	0	0	0	0	0	0	0	0	0
R	$\mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{Z}_2 \times \mathbb{Z}_8$	$\mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^3)}$	$\mathbb{Z}_2 \times \frac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$	$\mathbb{Z}_2 \times \mathbb{Z}_9$	$\mathbb{Z}_2 \times \frac{\mathbb{Z}_3[x]}{(x^2)}$	$\mathbb{Z}_3 \times \mathbb{Z}_9$	$\mathbb{Z}_3 \times \frac{\mathbb{Z}_3[x]}{(x^2)}$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3$
$\|R\|$	12	16	16	16	18	18	27	27	8	12
$\|Spec(R)\|$	2	2	2	2	2	2	2	2	3	3
$char(R)$	6	8	2	4	18	6	9	3	2	6
$\|V(\Gamma(R))\|$	7	11	11	11	11	11	14	14	6	9
$\gamma(\Gamma(R))$	0	0	0	0	0	0	0	0	0	0

Table 3. Non-local rings with planar zero-divisor graphs.

R	$\mathbb{F}_4 \times \mathbb{F}_4$	$\mathbb{F}_4 \times \mathbb{Z}_5$	$\mathbb{Z}_5 \times \mathbb{Z}_5$	$\mathbb{Z}_4 \times \mathbb{F}_4$	$\mathbb{F}_4 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{F}_4 \times \mathbb{Z}_7$	$\mathbb{Z}_4 \times \mathbb{Z}_4$	$\mathbb{Z}_4 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\frac{\mathbb{Z}_2[x]}{(x^2)} \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x,y]}{(x^2,xy,y^2)}$
$\|R\|$	16	20	25	16	16	28	16	16	16	16
$\|Spec(R)\|$	2	2	2	2	2	2	2	2	2	2
$char(R)$	2	10	5	4	2	14	4	4	2	2
$\|V(\Gamma(R))\|$	6	7	8	9	9	9	11	11	11	11
$\gamma(\Gamma(R))$	1	1	1	1	1	1	1	1	1	1
R	$\mathbb{Z}_2 \times \frac{\mathbb{Z}_4[x]}{(2x,x^2)}$	$\mathbb{Z}_4 \times \mathbb{Z}_5$	$\mathbb{Z}_5 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{Z}_3 \times \mathbb{Z}_8$	$\mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^3)}$	$\mathbb{Z}_3 \times \frac{\mathbb{Z}_4[x]}{(x^2-2,x^3)}$	$\mathbb{Z}_4 \times \mathbb{Z}_7$	$\mathbb{Z}_7 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{Z}_2 \times \frac{\mathbb{F}_4[x]}{(x^2)}$	$\mathbb{Z}_2 \times \frac{\mathbb{Z}_4[x]}{(x^2+x+1)}$
$\|R\|$	16	20	20	24	24	24	28	28	32	32
$\|Spec(R)\|$	2	2	2	2	2	2	2	2	2	2
$char(R)$	4	20	10	24	6	12	28	14	2	4
$\|V(\Gamma(R))\|$	11	11	11	15	15	15	15	15	19	19
$\gamma(\Gamma(R))$	1	1	1	1	1	1	1	1	1	1
R	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{F}_4$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_4$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$	$\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_5$	$\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{F}_4$	$\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_7$	$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$	
$\|R\|$	16	16	16	18	20	24	27	28	16	
$\|Spec(R)\|$	3	3	3	3	3	3	3	3	4	
$char(R)$	2	4	2	6	10	6	3	14	2	
$\|V(\Gamma(R))\|$	12	13	13	13	15	17	18	21	14	
$\gamma(\Gamma(R))$	1	1	1	1	1	1	1	1	1	

Table 4. Non-local rings with toroidal zero-divisor graphs.

References

[1] M. Atiyah and I. MacDonald, *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass., 1969.

[2] S. Akbari, H.R. Maimani and S. Yassemi, *When a zero-divisor graph is planar or a complete r-partite graph*, J. Algebra **270** (2003), 169–180.

[3] S. Akbari and A. Mohammadian, *On the zero-divisor graph of a commutative ring*, J. Algebra **274** (2004), 847–855.

[4] D.F. Anderson, A. Frazier, A. Lauve and P.S. Livingston, *The zero-divisor graph of a commutative ring, II*, Lecture Notes in Pure and Appl. Math. **220** (2001), 61–72.

[5] D.F. Anderson, R. Levy, J. Shapiro, *Zero-divisor graphs, von Neumann regular rings, and Boolean algebras*, J. Pure Appl. Algebra 180 (2003), no. 3, 221–241.

[6] D.F. Anderson and P.S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), 434–447.

[7] R. Belshoff and J. Chapman, *Planar Zero-Divisor Graphs*, J. Algebra **316** (2007), 471–480.

[8] F. DeMeyer and K. Schneider, *Automorphisms and zero-divisor graphs of commutative rings*, Commutative rings, 25–37, Nova Sci. Publ., Hauppauge, NY, 2002.

[9] C. Godsil and G. Royle, *Algebraic graph theory*, Graduate Texts in Mathematics, 207. Springer-Verlag, New York, 2001.

[10] J.L. Gross and T.W. Tucker, *Topological Graph Theory*, Wiley Interscience, New York, 1987.

[11] F. Harary, *Graph Theory*, Addison-Wesley Publishing Co., Reading Mass., 1972.

[12] A. Lauve, *Zero-divisor graphs of finite commutative rings*, Honors Senior Thesis, The University of Oklahoma, Norman, OK, April, 1999.

[13] P.S. Livingston, *Structure in zero-divisor graphs of commutative rings*, Masters Thesis, The University of Tennessee, Knoxville, TN, December 1997.

[14] J. Mayer, *Le problème des régions voisines sur les surfaces closes orientables*, J. Combinatorial Theory 6 (1969), 177–195.

[15] S.B. Mulay, *Cycles and symmetries of zero-divisors*, Communications in Algebra **30**(7) (2002), 3533–3558.

[16] S.P. Redmond, *On zero-divisor graphs of small finite commutative rings*, Discrete Math. **307** (2007), 1155–1166.

[17] G. Ringel and J.W.T. Youngs, *Solution of the Heawood map-coloring problem*, Proc. Nat. Acad. Sci. USA **60** (1968), 438–445.

[18] N. Smith, *Planar Zero-Divisor Graphs*, International Journal of Commutative Rings **2(4)** (2003), 177–188.

[19] N. Smith, *Infinite Planar Zero-Divisor Graphs*, Communications in Algebra **35** (2007), 171–180.

[20] H.-J. Wang, *Zero-divisor graphs of genus one*, J. Algebra **304** (2006), 666-678.

[21] C. Wickham, *Classification of Rings with Genus One Zero-Divisor Graphs*, Preprint.